

Product datasheet

Anti-AKR1C3 antibody ab84327

7 References 3 Images

Overview

Product name	Anti-AKR1C3 antibody
Description	Rabbit polyclonal to AKR1C3
Host species	Rabbit
Tested applications	Suitable for: WB, IHC-P, ICC/IF
Species reactivity	Reacts with: Cow, Human
Immunogen	A 15 amino acid peptide corresponding to the C terminal residues of human AKR1C3 (P42330).
Positive control	Human fetal liver lysates; human lung tissue; HeLa cells.

Properties

Form	Lyophilised:Reconstitute with 200ul distilled sterile water. Please note that if you receive this product in liquid form it has already been reconstituted as described and no further reconstitution is necessary.
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
Storage buffer	Preservative: 0.02% Sodium Azide Constituents: 2% BSA
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab84327** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
WB		1/500 - 1/1000. Predicted molecular weight: 36 kDa.
IHC-P		1/100 - 1/500.

Application	Abreviews	Notes
ICC/IF		1/50 - 1/200.

Target

Function	Catalyzes the conversion of aldehydes and ketones to alcohols. Catalyzes the reduction of prostaglandin (PG) D2, PGH2 and phenanthrenequinone (PQ) and the oxidation of 9-alpha,11-beta-PGF2 to PGD2. Functions as a bi-directional 3-alpha-, 17-beta- and 20-alpha HSD. Can interconvert active androgens, estrogens and progestins with their cognate inactive metabolites. Preferentially transforms androstenedione (4-dione) to testosterone.
Tissue specificity	Expressed in many tissues including adrenal gland, brain, kidney, liver, lung, mammary gland, placenta, small intestine, colon, spleen, prostate and testis. The dominant HSD in prostate and mammary gland. In the prostate, higher levels in epithelial cells than in stromal cells. In the brain, expressed in medulla, spinal cord, frontotemporal lobes, thalamus, subthalamic nuclei and amygdala. Weaker expression in the hippocampus, substantia nigra and caudate.
Sequence similarities	Belongs to the aldo/keto reductase family.
Cellular localization	Cytoplasm.


Images




Anti-AKR1C3 antibody (ab84327) at 1/500 dilution + human fetal liver lysate

Predicted band size: 36 kDa

Observed band size: 36 kDa



ab84327, at a 1/50 dilution, staining AKR1C3 in HeLa cells by Immunofluorescence.



ab84327, at a 1/100 dilution, staining AKR1C3 in formalin fixed, paraffin embedded human lung tissue by Immunohistochemistry. Note cytoplasmic staining.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AKR1C3 antibody (ab84327)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors