

Product datasheet

Anti-Apolipoprotein E antibody (Biotin) ab24274

1 References 2 Images

Overview

Product name	Anti-Apolipoprotein E antibody (Biotin)
Description	Goat polyclonal to Apolipoprotein E (Biotin)
Host species	Goat
Conjugation	Biotin
Tested applications	Suitable for: ICC/IF, ELISA, WB
Species reactivity	Reacts with: Human
Immunogen	Purified human Apolipoprotein E from human plasma.
Positive control	Liver (Human), Brain (Human), Spleen (Human), Ovary (Human), Kidney (Human) This antibody gave a positive result when used in the following methanol fixed cell lines: HepG2.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.02% Sodium Azide Constituents: 75mM PBS, 75mM Sodium chloride, 1mM EDTA. pH 7.2
Purity	Immunogen affinity purified
Purification notes	Purified by human apo E-Sepharose chromatography, then labeled with biotin.
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab24274** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		Use a concentration of 5 µg/ml.

Application	Abreviews	Notes
ELISA		1/5000 - 1/40000.
WB		1/5000 - 1/40000. Predicted molecular weight: 36 kDa. Dilution optimised using Chromogenic detection.

Target

Function	Mediates the binding, internalization, and catabolism of lipoprotein particles. It can serve as a ligand for the LDL (apo B/E) receptor and for the specific apo-E receptor (chylomicron remnant) of hepatic tissues.
Tissue specificity	Occurs in all lipoprotein fractions in plasma. It constitutes 10-20% of very low density lipoproteins (VLDL) and 1-2% of high density lipoproteins (HDL). APOE is produced in most organs. Significant quantities are produced in liver, brain, spleen, lung, adrenal, ovary, kidney and muscle.
Involvement in disease	<p>Defects in APOE are a cause of hyperlipoproteinemia type 3 (HLPP3) [MIM:107741]; also known as familial dysbetalipoproteinemia. Individuals with HLPP3 are clinically characterized by xanthomas, yellowish lipid deposits in the palmar crease, or less specific on tendons and on elbows. The disorder rarely manifests before the third decade in men. In women, it is usually expressed only after the menopause. The vast majority of the patients are homozygous for APOE*2 alleles. More severe cases of HLPP3 have also been observed in individuals heterozygous for rare APOE variants. The influence of APOE on lipid levels is often suggested to have major implications for the risk of coronary artery disease (CAD). Individuals carrying the common APOE*4 variant are at higher risk of CAD.</p> <p>Genetic variations in APOE are associated with Alzheimer disease type 2 (AD2) [MIM:104310]. It is a late-onset neurodegenerative disorder characterized by progressive dementia, loss of cognitive abilities, and deposition of fibrillar amyloid proteins as intraneuronal neurofibrillary tangles, extracellular amyloid plaques and vascular amyloid deposits. The major constituent of these plaques is the neurotoxic amyloid-beta-APP 40-42 peptide (s), derived proteolytically from the transmembrane precursor protein APP by sequential secretase processing. The cytotoxic C-terminal fragments (CTFs) and the caspase-cleaved products such as C31 derived from APP, are also implicated in neuronal death. Note=The APOE*4 allele is genetically associated with the common late onset familial and sporadic forms of Alzheimer disease. Risk for AD increased from 20% to 90% and mean age at onset decreased from 84 to 68 years with increasing number of APOE*4 alleles in 42 families with late onset AD. Thus APOE*4 gene dose is a major risk factor for late onset AD and, in these families, homozygosity for APOE*4 was virtually sufficient to cause AD by age 80. The mechanism by which APOE*4 participates in pathogenesis is not known.</p> <p>Defects in APOE are a cause of sea-blue histiocyte disease (SBHD) [MIM:269600]; also known as sea-blue histiocytosis. This disorder is characterized by splenomegaly, mild thrombocytopenia and, in the bone marrow, numerous histiocytes containing cytoplasmic granules which stain bright blue with the usual hematologic stains. The syndrome is the consequence of an inherited metabolic defect analogous to Gaucher disease and other sphingolipidoses.</p> <p>Defects in APOE are a cause of lipoprotein glomerulopathy (LPG) [MIM:611771]. LPG is an uncommon kidney disease characterized by proteinuria, progressive kidney failure, and distinctive lipoprotein thrombi in glomerular capillaries. It mainly affects people of Japanese and Chinese origin. The disorder has rarely been described in Caucasians.</p>
Sequence similarities	Belongs to the apolipoprotein A1/A4/E family.

Post-translational modifications

Synthesized with the sialic acid attached by O-glycosidic linkage and is subsequently desialylated in plasma. O-glycosylated with core 1 or possibly core 8 glycans. Thr-307 is a minor glycosylation site compared to Ser-308.


Glycated in plasma VLDL of normal subjects, and of hyperglycemic diabetic patients at a higher level (2-3 fold).

Phosphorylation sites are present in the extracellular medium.

Cellular localization

Secreted.


Images



Copyright (c) 2013 abcam plc

Immunocytochemistry/ Immunofluorescence - Anti-Apolipoprotein E antibody (Biotin) (ab24274)

ICC/IF image of ab24274 stained HepG2 cells. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal donkey serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody ab24274 at 5µg/ml overnight at +4°C. The secondary antibody (green) was DyLight® 488 donkey anti- goat ([ab96931](#)) IgG (H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Western blot - Anti-Apolipoprotein E antibody
(Biotin) (ab24274)

All lanes : Anti-Apolipoprotein E antibody
(Biotin) (ab24274) at 1 µg/ml

Lane 1 : Human liver tissue lysate - total
protein (ab29889)

Lane 2 : Human brain tissue lysate - total
protein (ab29466)

Lane 3 : Human spleen tissue lysate - total
protein (ab29699)

Lane 4 : Human ovary tissue lysate - total
protein (ab30222)

Lane 5 : Human kidney tissue lysate - total
protein (ab30203)

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Rabbit polyclonal to Goat IgG -
H&L - Pre-Adsorbed (HRP) (ab65486) at
1/3000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 36 kDa

Observed band size: 36 kDa

Exposure time: 20 minutes

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors