

Product datasheet

BMP7 overexpression 293T lysate (whole cell) ab94187

2 Images

Overview

|                            |   |
|----------------------------|---|
| <b>Product name</b>        | BMP7 overexpression 293T lysate (whole cell)  |
| <b>Description</b>         | BMP7 overexpression lysate  |
| <b>General notes</b>       | ab94187 is a 293T cell transfected lysate in which Human BMP7 has been transiently over-expressed using a pCMV-BMP7 plasmid. The lysate is provided in 1X Sample Buffer. <b>Note:</b> For more details about how the transfected lysate was prepared <a href="#">view the preparation notes</a> . |
| <b>Tested applications</b> | <b>Suitable for:</b> WB   |

Properties

|                             |  |
|-----------------------------|--|
| <b>Form</b>                 | Liquid   |
| <b>Storage instructions</b> | Shipped on dry ice. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.  |
| <b>Storage buffer</b>       | pH: 6.80<br>Constituent: 100% 1x Sample Buffer   |
| <b>Purity</b>               | Whole Cell Lysate  |
| <b>Background</b>           | Function: Induces cartilage and bone formation. May be the osteoinductive factor responsible for the phenomenon of epithelial osteogenesis. Plays a role in calcium regulation and bone homeostasis. Tissue specificity: Expressed in the kidney and bladder. Lower levels seen in the brain. Similarity: Belongs to the TGF-beta family. Developmental stage: Expressed in the developing eye, brain and ear during embryogenesis. PTM: Several N-termini starting at positions 293, 300, 315 and 316 have been identified by direct sequencing resulting in secretion of different mature forms (PubMed:17977014). |


Applications

Our [Abpromise guarantee](#) covers the use of **ab94187** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes                               |
|-------------|-----------|-------------------------------------|
| WB          |           | Use at an assay dependent dilution. |

Images



ab94187 at 15 µg/lane on an SDS-PAGE gel



**All lanes** : Anti-BMP7 antibody (ab54904) at  
1/500 dilution

**Lane 1** : BMP7 overexpression 293T lysate  
(whole cell) (ab94187)

**Lane 2** : 293T non-transfected Lysate

Lysates/proteins at 25 µg per lane.

**Secondary**

Goat Anti-mouse IgG (H and L) HRP  
conjugated at 1/2500 dilution

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

**Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

**Terms and conditions**

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors