

Product datasheet

Anti-C4c+C4b antibody (FITC) ab4216

Overview

Product name	Anti-C4c+C4b antibody (FITC)
Description	Rabbit polyclonal to C4c+C4b (FITC)
Host species	Rabbit
Conjugation	FITC. Ex: 493nm, Em: 528nm
Specificity	This antibody reacts with human C4, C4b and C4c but does not react with the C4d epitope.
Tested applications	Suitable for: IHC-Fr, IHC-P
Species reactivity	Reacts with: Human
Immunogen	C4c complement isolated from complement activated human serum.
General notes	Fluorescein isothiocyanate (FITC) isomer 1.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C.
Storage buffer	Preservative: 0.05% Sodium Azide Constituents: 1% BSA, PBS
Purity	IgG fraction
Purification notes	Traces of contaminating antibodies have been removed by solid phase absorption with human plasma proteins.
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab4216** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
-------------	-----------	-------

Application	Abreviews	Notes
IHC-Fr		1/50 - 1/100. Incubate for 30 - 60 min at room temperature. Optimal dilutions / concentrations should be determined by the end user.
IHC-P		1/50 - 1/100. Incubate for 30 - 60 min at room temperature. Optimal dilutions / concentrations should be determined by the end user.

Target

Relevance

C4 plays a central role in the activation of the classical pathway of the complement system. Human complement component C4 is polymorphic at two loci, C4A and C4B. The C4A alleles carry the Rodgers (Rg) while the C4B alleles carry the Chido (Ch) blood group antigens. C4 (either allele A or B) protein is expressed as a single chain precursor which is proteolytically cleaved into a trimer of alpha, beta, and gamma chains prior to secretion. The trimer provides a surface for interaction between the antigen-antibody complex and other complement components. During activation, the alpha chain is cleaved by C1 into C4a and C4b, and C4b stays linked to the beta and gamma chains. Further degradation of C4b by C1 into the inactive fragments C4c and C4d blocks the generation of C3 convertase.

Cellular localization

Secreted

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors