

Product datasheet

# Anti-C5R1 antibody ab59390

[1 Abreviews](#) [1 References](#) [2 Images](#)

## Overview

<b>Product name</b>	Anti-C5R1 antibody
<b>Description</b>	Rabbit polyclonal to C5R1
<b>Specificity</b>	This antibody detects endogenous levels of total C5R1 protein.
<b>Tested applications</b>	<b>Suitable for:</b> ELISA, WB, IHC-P
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Human
<b>Immunogen</b>	Synthesized non-phosphopeptide derived from human C5R1 around the phosphorylation site of serine 338.
<b>Positive control</b>	WB: Extracts from HeLa cells treated with PMA (125ng/ml, 30mins).

## Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
<b>Storage buffer</b>	Preservative: 0.02% Sodium Azide Constituents: 50% Glycerol, PBS, 150mM Sodium chloride, pH 7.4
<b>Purity</b>	Immunogen affinity purified
<b>Purification notes</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG

## Applications

Our [Abpromise guarantee](#) covers the use of **ab59390** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ELISA		
WB		
IHC-P		

## Application notes

ELISA: 1/5000.

IHC-P: Use at a concentration of 1 µg/ml.

WB: 1/500 - 1/1000. Detects a band of approximately 46 kDa (predicted molecular weight: 39 kDa).

Lysis buffer: 50mM Tris(pH7.4),150mM NaCl,1% Triton X-100,1% sodium deoxycholate,0.1% SDS and sodium orthovanadate,sodium fluoride,EDTA ,leupeptin.

Blocking buffer: 3%BSA, 2 hours. Dilute the primary antibody in blocking buffer.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

## Target

### Function

Receptor for the chemotactic and inflammatory peptide anaphylatoxin C5a. This receptor stimulates chemotaxis, granule enzyme release and superoxide anion production.

### Sequence similarities

Belongs to the G-protein coupled receptor 1 family.


### Post-translational modifications

Sulfation plays a critical role in the association of the receptor with C5a, but no significant role in the ability of the receptor to transduce a signal and mobilize calcium in response to a small peptide agonist.

### Cellular localization

Cell membrane.

## Anti-C5R1 antibody images




**All lanes** : Anti-C5R1 antibody (ab59390) at 1/500 dilution

**Lane 1** : Extracts from HeLa cells, treated with PMA (125ng/ml, 30mins).

**Lane 2** : Extracts from HeLa cells, treated with PMA (125ng/ml, 30mins) plus peptide.

**Predicted band size** : 39 kDa

**Observed band size** : 46 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)-C5R1 antibody(ab59390)

ab59390 staining C5R1 in human lung.

Left panel: with primary antibody at 1 ug/ml.

Right panel: isotype control.

Sections were stained using an automated system (DAKO Autostainer Plus), at room temperature: sections were rehydrated and antigen retrieved with the Dako 3 in 1 AR buffers EDTA pH 9.0. Slides were peroxidase blocked in 3% H<sub>2</sub>O<sub>2</sub> in methanol for 10 mins. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 min and detected with Dako envision flex amplification kit for 30 minutes. Colorimetric detection was completed with Diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that for manual staining we recommend to optimize the primary antibody concentration and incubation time (overnight incubation), and amplification may be required.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

### **Terms and conditions**

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors