

## Product datasheet

# Anti-CCR4 antibody ab1669

★★★★☆ 4 Abreviews 10 References 8 Images

### Overview

<b>Product name</b>	Anti-CCR4 antibody
<b>Description</b>	Goat polyclonal to CCR4
<b>Host species</b>	Goat
<b>Specificity</b>	Peptide sequence is < 50 % identical to other human chemokine receptors in this region.
<b>Tested applications</b>	<b>Suitable for:</b> Flow Cyt, ICC/IF, ELISA, ICC, IHC-P, WB, IHC-Fr
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Human, Cynomolgus monkey
<b>Immunogen</b>	Synthetic peptide: SNYYLYESIPKPCTKEGIKAFGE , corresponding to amino acids 17-39 of Human CCR4 (extracellular domain). <a href="#">Run BLAST with</a> <a href="#">Run BLAST with</a>
<b>Positive control</b>	Human peripheral blood cells or paraffin sections of spleen. IHC-P:FFPE mouse lymph node normal. IHC-P:FFPE human tonsil normal. IHC-P:FFPE rat spleen normal.

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
<b>Storage buffer</b>	PBS, 1mg/ml BSA, 0.1% sodium azide, pH7.2
<b>Purity</b>	Immunogen affinity purified
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG

### Applications

Our [Abpromise guarantee](#) covers the use of **ab1669** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt	★★★★☆	Use at an assay dependent concentration. <a href="#">ab37373</a> - Goat polyclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF		Use at an assay dependent concentration. Fix cells with 4% PFA (see Ritter et al).
ELISA		1/100000.
ICC		1/300.
IHC-P		1/300.
WB		1/1000. Detects a band of approximately 55 kDa (predicted molecular weight: 42 kDa).
IHC-Fr		Use at an assay dependent concentration. Fix in acetone at -20°C for 5 minutes. See Heller et al.

## Target

<b>Function</b>	High affinity receptor for the C-C type chemokines CCL17/TARC and CCL22/MDC. The activity of this receptor is mediated by G(i) proteins which activate a phosphatidylinositol-calcium second messenger system. Can function as a chemoattractant homing receptor on circulating memory lymphocytes and as a coreceptor for some primary HIV-2 isolates. In the CNS, could mediate hippocampal-neuron survival.
<b>Tissue specificity</b>	Predominantly expressed in the thymus, in peripheral blood leukocytes, including T-cells, mostly CD4+ cells, and basophils, and in platelets; at lower levels, in the spleen and in monocytes. Detected also in macrophages, IL-2-activated natural killer cells and skin-homing memory T-cells, mostly the ones expressing the cutaneous lymphocyte antigen (CLA). Expressed in brain microvascular and coronary artery endothelial cells.
<b>Sequence similarities</b>	Belongs to the G-protein coupled receptor 1 family.
<b>Post-translational modifications</b>	In natural killer cells, CCL22 binding induces phosphorylation on yet undefined Ser/Thr residues, most probably by beta-adrenergic receptor kinases 1 and 2.
<b>Cellular localization</b>	Cell membrane.

## Images




Flow Cytometry - Anti-CCR4 antibody (ab1669)

This image is courtesy of an anonymous Abreview

ab1669 staining CCR4 in Mouse whole blood by Flow Cytometry. Red blood cells were lysed in PBS + 1% BSA and 0.01% sodium azide and fixed in paraformaldehyde. The sample was incubated with the primary antibody (1/100 in PBS + 1% BSA and 0.01% sodium azide) for 1 hour at 4°C. A biotin-conjugated Donkey anti-goat IgG polyclonal (1/200) was used as the secondary antibody.


**Gating Strategy:** Live Lymphocytes.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CCR4 antibody (ab1669)

IHC image of CD3 staining in a formalin fixed, paraffin embedded normal rat spleen tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab1669 at 1/100 dilution for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.


For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Immunohistochemistry (Frozen sections) - Anti-CCR4 antibody (ab16669)

Frozen sections of human skin tissue stained for CCR4 using ab16669 in immunohistochemical analysis (green). CD31(red) and nuclei (blue) are also shown. The right hand panel is a merged image.

Image courtesy of PMID 25915746 (PLoS One 2015 Apr 27;10(4)).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CCR4 antibody (ab16669)

IHC image of CD3 staining in a formalin fixed, paraffin embedded normal human tonsil tissue section\*, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab16669 at 1/100 dilution for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.


\*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CCR4 antibody (ab1669)


IHC image of CD3 staining in TISSUE formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol B. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab1669 at 1/100 dilution for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.




- CCR4 antibody (ab1669)

Immunohistochemistry using ab1669 on a section of human spleen.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CCR4 antibody (ab1669)

IHC image of CCR4 staining in human spleen formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab1669, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Anti-CCR4 antibody (ab1669) at 1 µg/ml +  
Human thymus tissue lysate - total protein  
(ab30146) at 10 µg

### Secondary

Rabbit polyclonal to Goat IgG - H&L - Pre-  
Adsorbed (HRP) at 1/3000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

**Predicted band size:** 42 kDa

**Observed band size:** 55 kDa

**Additional bands at:** 100 kDa, 37 kDa. We  
are unsure as to the identity of these extra  
bands.

**Exposure time:** 12 minutes

CCR4 contains a number of potential  
glycosylation sites (SwissProt) which may  
explain its migration at a higher molecular  
weight than predicted.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors