

## Product datasheet

# Anti-CTCF antibody - ChIP Grade ab70303

★★★★★ 7 Abreviews 26 References 5 Images

### Overview

<b>Product name</b>	Anti-CTCF antibody - ChIP Grade
<b>Description</b>	Rabbit polyclonal to CTCF - ChIP Grade
<b>Tested applications</b>	<b>Suitable for:</b> WB, IP, ChIP, IHC-P, EMSA
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Human <b>Predicted to work with:</b> Rat, Cow, Pig, Chimpanzee, Rhesus monkey, Gorilla, Orangutan, Elephant
<b>Immunogen</b>	Synthetic peptide corresponding to a region between residues 650 and 700 of human CTCF (Peptide available as <a href="#">ab115042</a> .)
<b>Positive control</b>	MCF7, 293T, NIH3T3 and HeLa whole cell lysates and Primary human lung fibroblasts.

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	pH: 7.80 Preservative: 0.09% Sodium azide Constituents: 1.815% Tris, 1.764% Sodium citrate, 0.021% PBS
<b>Purity</b>	Immunogen affinity purified
<b>Purification notes</b>	ab70303 was affinity purified using an epitope specific to CTCF immobilized on solid support.
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG

### Applications

Our [Abpromise guarantee](#) covers the use of **ab70303** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	★★★★★	1/2500 - 1/10000. Detects a band of approximately 83 kDa (predicted molecular weight: 83 kDa).

Application	Abreviews	Notes
IP		Use at 2-5 µg/mg of lysate.
ChIP	★★★★★	Use at an assay dependent concentration.
IHC-P		1/100 - 1/500. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
EMSA		Use at an assay dependent concentration. PubMed: 22210889

## Target

### Function

Chromatin binding factor that binds to DNA sequence specific sites. Involved in transcriptional regulation by binding to chromatin insulators and preventing interaction between promoter and nearby enhancers and silencers. Acts as transcriptional repressor binding to promoters of vertebrate MYC gene and BAG1 gene. Also binds to the PLK and PIM1 promoters. Acts as a transcriptional activator of APP. Regulates APOA1/C3/A4/A5 gene cluster and controls MHC class II gene expression. Plays an essential role in oocyte and preimplantation embryo development by activating or repressing transcription. Seems to act as tumor suppressor. Plays a critical role in the epigenetic regulation. Participates to the allele-specific gene expression at the imprinted IGF2/H19 gene locus. On the maternal allele, binding within the H19 imprinting control region (ICR) mediates maternally inherited higher-order chromatin conformation to restrict enhancer access to IGF2. Plays a critical role in gene silencing over considerable distances in the genome. Preferentially interacts with unmethylated DNA, preventing spreading of CpG methylation and maintaining methylation-free zones. Inversely, binding to target sites is prevented by CpG methylation. Plays an important role in chromatin remodeling. Can dimerize when it is bound to different DNA sequences, mediating long-range chromatin looping. Mediates interchromosomal association between IGF2/H19 and WSB1/NF1 and may direct distant DNA segments to a common transcription factory. Causes local loss of histone acetylation and gain of histone methylation in the beta-globin locus, without affecting transcription. When bound to chromatin, it provides an anchor point for nucleosomes positioning. Seems to be essential for homologous X-chromosome pairing. May participate with Tsix in establishing a regulatable epigenetic switch for X chromosome inactivation. May play a role in preventing the propagation of stable methylation at the escape genes from X-inactivation. Involved in sister chromatid cohesion. Associates with both centromeres and chromosomal arms during metaphase and required for cohesin localization to CTCF sites. Regulates asynchronous replication of IGF2/H19.

### Tissue specificity

Ubiquitous. Absent in primary spermatocytes.

### Sequence similarities

Belongs to the CTCF zinc-finger protein family.  
Contains 11 C2H2-type zinc fingers.

### Domain

The 11 zinc fingers are highly conserved among vertebrates, exhibiting almost identical amino acid sequences. Different subsets or combination of individual zinc fingers gives the ability to CTCF to recognize multiple DNA target sites.

### Post-translational modifications


Sumoylated on Lys-74 and Lys-689; sumoylation of CTCF contributes to the repressive function of CTCF on the MYC P2 promoter.

### Cellular localization

Nucleus > nucleoplasm. Chromosome. Chromosome > centromere. May translocate to the nucleolus upon cell differentiation. Associates with both centromeres and chromosomal arms

during metaphase. Associates with the H19 ICR in mitotic chromosomes. May be preferentially excluded from heterochromatin during interphase.

## Images




Western blot - Anti-CTCF antibody - ChIP Grade  
(ab70303)

**Predicted band size :** 83 kDa

Samples: Whole cell lysate (50 µg) from CT26.WT, 4T1, Renca, and TCMK-1 cells. ab70303 used at a dilution of 0.2 µg/ml.


**Detection:** Chemiluminescence with an exposure time of 30 seconds.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CTCF antibody - ChIP Grade (ab70303)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human ovarian carcinoma (left) and mouse renal cell carcinoma (right) tissues labelling CTCF with ab70303 at 1/100 (2µg/ml).

Detection: DAB.



Western blot - Anti-CTCF antibody - ChIP Grade (ab70303)

**All lanes :** Anti-CTCF antibody - ChIP Grade (ab70303) at 0.03 µg/ml

**Lane 1 :** MCF7 whole cell lysate at 50 µg

**Lane 2 :** MCF7 whole cell lysate at 15 µg

**Lane 3 :** MCF7 whole cell lysate at 5 µg

**Lane 4 :** 293T whole cell lysate at 50 µg


**Lane 5 :** NIH3T3 whole cell lysate at 50 µg

**Lane 6 :** HeLa whole cell lysate at 50 µg

**Predicted band size :** 83 kDa


**Observed band size :** 83 kDa

**Exposure time :** 30 seconds



Immunoprecipitation - Anti-CTCF antibody - ChIP Grade (ab70303)

Detection of Human CTCF by Western Blot of Immunoprecipitate. Lane 1: MCF7 whole cell lysates immunoprecipitated using [ab70302](#) at 3µg/mg lysate (1 mg/IP; 1/4 of IP loaded/lane). Lane 2: MCF7 whole cell lysates immunoprecipitated using [ab70303](#) at 3µg/mg lysate (1 mg/IP; 1/4 of IP loaded/lane). Lane 3: MCF7 whole cell lysates immunoprecipitated using [ab70304](#) at 3µg/mg lysate (1 mg/IP; 1/4 of IP loaded/lane). Lane 4: control IgG was used for immunoprecipitation. All lanes: [ab70303](#) at 0.03µg/ml. Detection: Chemiluminescence with exposure times of 10 seconds.



ChIP - Anti-CTCF antibody - ChIP Grade (ab70303)

Analysis of Human CTCF by Chromatin Immunoprecipitation (ChIP). Primary human fibroblasts from lung ( $5 \times 10^7$  cells) were cross-linked with formaldehyde, sonicated, and immunoprecipitated with 20µg [ab70303](#). The resulting ChIP DNA was quantified using real-time PCR with primers against the IGF2-H19 ICR or a control region in the HoxA cluster.

## **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors