

## Product datasheet

# Anti-CaMKI antibody [EPR2217Y] ab68234

Recombinant RabMAb

★★★★☆ 2 Abreviews 6 References 2 Images

### Overview

<b>Product name</b>	Anti-CaMKI antibody [EPR2217Y]
<b>Description</b>	Rabbit monoclonal [EPR2217Y] to CaMKI
<b>Tested applications</b>	<b>Suitable for:</b> WB, IHC-P, ICC <b>Unsuitable for:</b> Flow Cyt or IP
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Rat, Human
<b>Immunogen</b>	A specific peptide corresponding to residues on the N-terminus of human CaMKI.
<b>Positive control</b>	SH-SY5Y lysate, human, mouse and rat brain lysate and human kidney tissue sections.
<b>General notes</b>	This product is a recombinant rabbit monoclonal antibody. Our RabMAb <sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
<b>Storage buffer</b>	PBS 49%,Sodium azide 0.01%,Glycerol 50%,BSA 0.05%
<b>Purity</b>	Tissue culture supernatant
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	EPR2217Y
<b>Isotype</b>	IgG

### Applications

Our [Abpromise guarantee](#) covers the use of **ab68234** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	★★★★☆	1/5000 - 1/10000. Detects a band of approximately 41 kDa (predicted molecular weight: 41 kDa).
IHC-P	★★★★☆	1/100 - 1/250.
ICC		1/100 - 1/250.

**Application notes** Is unsuitable for Flow Cyt or IP.

## Target

### Function

Calcium/calmodulin-dependent protein kinase that operates in the calcium-triggered CaMKK-CaMK1 signaling cascade and, upon calcium influx, regulates transcription activators activity, cell cycle, hormone production, cell differentiation, actin filament organization and neurite outgrowth. Recognizes the substrate consensus sequence [MVLIF]-x-R-x(2)-[ST]-x(3)-[MVLIF]. Regulates axonal extension and growth cone motility in hippocampal and cerebellar nerve cells. Upon NMDA receptor-mediated Ca(2+) elevation, promotes dendritic growth in hippocampal neurons and is essential in synapses for full long-term potentiation (LTP) and ERK2-dependent translational activation. Downstream of NMDA receptors, promotes the formation of spines and synapses in hippocampal neurons by phosphorylating ARHGEF7/BETAPIX on 'Ser-694', which results in the enhancement of ARHGEF7 activity and activation of RAC1. Promotes neuronal differentiation and neurite outgrowth by activation and phosphorylation of MARK2 on 'Ser-91', 'Ser-92', 'Ser-93' and 'Ser-294'. Promotes nuclear export of HDAC5 and binding to 14-3-3 by phosphorylation of 'Ser-259' and 'Ser-498' in the regulation of muscle cell differentiation. Regulates NUMB-mediated endocytosis by phosphorylation of NUMB on 'Ser-276' and 'Ser-295'. Involved in the regulation of basal and estrogen-stimulated migration of medulloblastoma cells through ARHGEF7/BETAPIX phosphorylation (By similarity). Is required for proper activation of cyclin-D1/CDK4 complex during G1 progression in diploid fibroblasts. Plays a role in K(+) and ANG2-mediated regulation of the aldosterone synthase (CYP11B2) to produce aldosterone in the adrenal cortex. Phosphorylates EIF4G3/eIF4GII. In vitro phosphorylates CREB1, ATF1, CFTR, MYL9 and SYN1/synapsin I.

### Tissue specificity

Widely expressed. Expressed in cells of the zona glomerulosa of the adrenal cortex.

### Sequence similarities

Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. CaMK subfamily.  
Contains 1 protein kinase domain.

### Domain

The autoinhibitory domain overlaps with the calmodulin binding region and interacts in the inactive folded state with the catalytic domain as a pseudosubstrate.


### Post-translational modifications

Phosphorylated by CaMKK1 and CaMKK2 on Thr-177.  
Polyubiquitinated by the E3 ubiquitin-protein ligase complex SCF(FBXL12), leading to proteasomal degradation.

### Cellular localization

Cytoplasm. Nucleus. Predominantly cytoplasmic.

## Images



Western blot - Anti-CaMKI antibody [EPR2217Y]  
(ab68234)

**All lanes :** Anti-CaMKI antibody [EPR2217Y]  
(ab68234) at 1/5000 dilution

**Lane 1 :** SH-SY5Y lysate

**Lane 2 :** human brain tissue lysate

**Lane 3 :** mouse brain tissue lysate

**Lane 4 :** rat brain tissue lysate


Lysates/proteins at 10 µg per lane.

**Secondary**

HRP labelled goat anti-rabbit at 1/2000  
dilution

**Predicted band size :** 41 kDa

**Observed band size :** 41 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-  
embedded sections) - Anti-CaMKI antibody  
[EPR2217Y] (ab68234)

ab68234 at 1/100-1/250 dilution staining  
CaMKI in human kidney tissue sections.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

**Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

**Terms and conditions**

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors