

Product datasheet

Anti-Caspase-3 antibody ab13847

KO VALIDATED

★★★★☆ 41 Abreviews 211 References 7 Images

Overview

Product name	Anti-Caspase-3 antibody
Description	Rabbit polyclonal to Caspase-3
Host species	Rabbit
Specificity	ab13847 recognizes a cleaved form of Caspase 3 (~17 kDa) after apoptosis has been induced in wildtype cells and not Caspase 3 knockout cells.
Tested applications	Suitable for: ICC/IF, IHC-P, IHC-Fr, WB, Flow Cyt, ICC, IHC (PFA fixed), IHC-FoFr, IHC - Wholemout
Species reactivity	Reacts with: Mouse, Rat, Human, Pig, Xenopus laevis, Drosophila melanogaster, Indian muntjac, Zebrafish, Rhesus monkey, Common marmoset, Schmidtea mediterranea, Salvelinus alpinus Predicted to work with: Dog, Chinese hamster
Immunogen	Synthetic peptide corresponding to Human Caspase-3 aa 150-250 (internal sequence) conjugated to keyhole limpet haemocyanin. Database link: P42574 (Peptide available as ab13848)
Positive control	This antibody gave a positive signal in WB with active Caspase 3 recombinant protein and pro-Caspase 3 recombinant protein. This antibody also gave a signal with HeLa staurosporine treated (2uM/4 hr) whole cell lysate.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS Note: Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab13847** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF	★★★★☆	Use a concentration of 5 µg/ml.
IHC-P	★★★★☆	1/50. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
IHC-Fr	★★★★★	Use at an assay dependent concentration.
WB	★★★★☆	1/500. Detects a band of approximately 17 kDa (predicted molecular weight: 17 kDa).
Flow Cyt	★★★★★	1/500. ab171870 - Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.
ICC	★★★★☆	Use at an assay dependent concentration.
IHC (PFA fixed)		1/300.
IHC-FoFr	★★★★★	1/300.
IHC - Wholemout	★★★★☆	1/500.

Target


Function	Involved in the activation cascade of caspases responsible for apoptosis execution. At the onset of apoptosis it proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at a '216-Asp-Gly-217' bond. Cleaves and activates sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Cleaves and activates caspase-6, -7 and -9. Involved in the cleavage of huntingtin.
Tissue specificity	Highly expressed in lung, spleen, heart, liver and kidney. Moderate levels in brain and skeletal muscle, and low in testis. Also found in many cell lines, highest expression in cells of the immune system.
Sequence similarities	Belongs to the peptidase C14A family.
Post-translational modifications	Cleavage by granzyme B, caspase-6, caspase-8 and caspase-10 generates the two active subunits. Additional processing of the propeptides is likely due to the autocatalytic activity of the activated protease. Active heterodimers between the small subunit of caspase-7 protease and the large subunit of caspase-3 also occur and vice versa. S-nitrosylated on its catalytic site cysteine in unstimulated human cell lines and denitrosylated upon activation of the Fas apoptotic pathway, associated with an increase in intracellular caspase activity. Fas therefore activates caspase-3 not only by inducing the cleavage of the

caspace zymogen to its active subunits, but also by stimulating the denitrosylation of its active site thiol.

Cellular localization

Cytoplasm.

Images



Lane 1: Wild-type HAP1 cell lysate + Staurosporine ([ab146588](#)) (1 μ M for 4h)


Lane 2: Wild-type HAP1 cell lysate

Lane 3: Caspase-3 knockout HAP1 cell lysate + Staurosporine ([ab146588](#)) (1 μ M for 4h)

Lane 4: Caspase-3 knockout HAP1 cell lysate

Lanes 1 - 4: Merged signal (red and green). Green - ab13847 observed at 17 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.


ab13847 was shown to recognise Caspase 3 when Caspase 3 knockout samples were used, along with additional cross-reactive bands. Wild-type and Caspase 3 knockout samples (\pm staurosporine treatment) were subjected to SDS-PAGE. ab13847 and [ab8245](#) (loading control to GAPDH) were diluted to 1/500 and 1/10000 respectively and incubated overnight at 4°C. Blots were developed with [Goat anti-Rabbit IgG H&L \(IRDye® 800CW\) preadsorbed \(ab216773\)](#) and [Goat anti-Mouse IgG H&L \(IRDye® 680RD\) preadsorbed \(ab216776\)](#) secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Caspase-3 antibody (ab13847)

This image is courtesy of an abreview from Zachary Yu-Ching Lin

IHC-P image of Caspase 3 staining with ab13847 on tissue sections from juvenile marmoset testis. The sections were subjected to heat-mediated antigen retrieval using Dako antigen retrieval solution. The sections were then blocked with 5% milk for 30 minutes at 25°C, before incubation with ab13847 (1/100 dilution) for 18 hours at 4°C. The secondary was an Alexa-Fluor 555 conjugated goat anti-rabbit polyclonal, used at a 1/500 dilution.




Flow Cytometry - Anti-Caspase-3 antibody (ab13847)

This image is courtesy of an anonymous Abreview

ab13847 staining active caspase 3 in Human Jurkat cells by Flow Cytometry. Cells were prepared in a phosphate buffered solution containing 0.1% sodium azide with FBS fixed with paraformaldehyde and permeabilized with Triton X-100 and NP40. The sample was incubated with the primary antibody (1/100 in wash buffer) for 24 hours at 4°C. A FITC-conjugated Goat anti-rabbit Ig (1/100) was used as the secondary antibody.


Gating Strategy: Isolate cell population from plot of SSC-A / FSA-A



Immunohistochemistry (Frozen sections) - Anti-Caspase-3 antibody (ab13847)

Image from PLoS One. 2015; 10(5): e0126688. Fig 2b, doi: 10.1371/journal.pone.0126688

5μm frozen sections of tumor tissue were fixed with 100% ice cold methanol for 10 minutes, then blocked in 5% normal goat serum in PBS (pH 7.4) for 1h. Sections were incubated with ab13847 (1:500) at 4°C overnight and for 1h with secondary antibodies at room temperature.



Western blot - Anti-Caspase-3 antibody (ab13847)

All lanes : Anti-Caspase-3 antibody (ab13847) at 1 μg/ml

Lane 1 : Human Caspase 3 (active) Recombinant Protein

Lane 2 : Human Pro Caspase 3 (inactive) Recombinant Protein

Lysates/proteins at 0.1 μg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/10000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 17 kDa

Observed band size: 17,32 kDa


Exposure time: 8 minutes

Caspase 3 exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce large

(17kDa) and small (12kDa) subunits. These subunits dimerize to form the active enzyme. ab13847 specifically detects the large active subunit (17kDa) and the inactive pro Caspase 3 (32 kDa).

This blot was produced using a 4-12% Bis-tris gel under the MES buffer system. The gel was run at 200V for 35 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 5% Bovine Serum Albumin before being incubated with ab13847 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution.


Secondary antibody - [Goat Anti-Rabbit IgG H&L \(HRP\) \(ab97051\) secondary antibody](#)



Immunohistochemistry (Frozen sections) - Anti-Caspase-3 antibody (ab13847)

This image is courtesy of an abreview submitted by Sophie Pezet, ESPCI, France

ab13847 was used in IHC of frozen sections from a rat brain with a kainite lesion. The non lesioned contralateral site serves as a negative control. The sections were fixed with paraformaldehyde. The tissue was perfused with 4% PFA and embedded in OCT compound and cut on the cryostat. The primary antibody was incubated for 12 hours at a dilution of 1/300. A biotin labelled secondary antibody was used at a dilution of 1/300.



Immunocytochemistry/ Immunofluorescence - Anti-Caspase-3 antibody (ab13847)

This image is courtesy of Roberto Giamb Bruno, Marilena Ciciarello and Patrizia Lavia

HeLa cells were fixed for 10 minutes at room temperature in 3.7% PFA and permeabilised in 0.1% Triton X-100/PBS then incubated with ab13847 (5µg/ml) for 1 hour at room temperature. The top panel shows control cells treated with DMSO. The bottom panel shows HeLa cells treated with 1 mM staurosporine (ab146588) for 4 hours to induce caspase-3 activation. ab13847 staining is shown in green and counterstaining with DAPI is shown in blue. 100x magnification.

The image shows the staining with ab13847 is very faint in the untreated control cultures, but very bright after activation of capsase-3 by treatment with the staurosporine . (N.B. in these cultures the nuclei are apoptotic).

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors