

Product datasheet

Anti-Cyclin D1 antibody [EPR2241] ab134175

Recombinant RabMAb

★★★★★ 7 Abreviews 60 References 15 Images

Overview

| | |
|----------------------------|--|
| Product name | Anti-Cyclin D1 antibody [EPR2241] |
| Description | Rabbit monoclonal [EPR2241] to Cyclin D1 |
| Specificity | The immunogen used for this product shares 66% homology with CCND2 (seven amino acid stretch with 100% homology). The cross-reactivity with this protein has not been confirmed experimentally. |
| Tested applications | Suitable for: WB, IP, ICC/IF, IHC-P |
| Species reactivity | Reacts with: Mouse, Rat, Human |
| Immunogen | Synthetic peptide (the amino acid sequence is considered to be commercially sensitive) corresponding to Human Cyclin D1 (C terminal). Database link: P24385 (Peptide available as ab188123) |
| Positive control | Human mantle cell lymphoma tissue; MCF7 cell lysate. ICC/IF: MCF7 cells treated with KN-93 IHC/P: Human esophagus (FFPE), Rat esophagus (FFPE) |
| General notes | This product is a recombinant rabbit monoclonal antibody. We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team. Produced using Abcam's RabMAb [®] technology. RabMAb [®] technology is covered by the following U.S. Patents, No. 5,675,063 and/or 7,429,487. A trial size is available to purchase for this antibody. |

Alternative versions available:

[Anti-Cyclin D1 antibody \(Alexa Fluor[®] 488\) \[EPR2241\] \(ab190194\)](#)

[Anti-Cyclin D1 antibody \(Alexa Fluor[®] 647\) \[EPR2241\] \(ab190563\)](#)

[Anti-Cyclin D1 antibody \(HRP\) \[EPR2241\] \(ab190564\)](#)

[Anti-Cyclin D1 antibody \(Alexa Fluor[®] 594\) \[EPR2241\] \(ab203446\)](#)

[Anti-Cyclin D1 antibody \(Alexa Fluor[®] 555\) \[EPR2241\] \(ab203448\)](#)

[Anti-Cyclin D1 antibody \(Alexa Fluor[®] 568\) \[EPR2241\] \(ab203449\)](#)

Properties

| | |
|--|---|
| Form | Liquid |
| Storage instructions | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C. |
| Dissociation constant (K_D) | $K_D = 2.40 \times 10^{-11} \text{ M}$ ^A Learn more about K_D |
| Storage buffer | pH: 7.40 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol, 0.05% BSA, 59% PBS |
| Purity | Immunogen affinity purified |
| Clonality | Monoclonal |
| Clone number | EPR2241 |
| Isotype | IgG |

Applications

Our [Abpromise guarantee](#) covers the use of **ab134175** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|-------------|-----------|--|
| WB | ★★★★★ | 1/10000 - 1/50000. Predicted molecular weight: 34 kDa. |
| IP | | 1/30. |
| ICC/IF | ★★★★★ | 1/50. |
| IHC-P | | 1/100 - 1/500. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See protocols (link: http://www.abcam.com/protocols/ihc-antigen-retrieval-protocol). |

Target

| | |
|-------------------------------|--|
| Function | Essential for the control of the cell cycle at the G1/S (start) transition. |
| Involvement in disease | Note=A chromosomal aberration involving CCND1 may be a cause of B-lymphocytic malignancy, particularly mantle-cell lymphoma (MCL). Translocation t(11;14)(q13;q32) with immunoglobulin gene regions. Activation of CCND1 may be oncogenic by directly altering progression through the cell cycle. Note=A chromosomal aberration involving CCND1 may be a cause of parathyroid adenomas. Translocation t(11;11)(q13;p15) with the parathyroid hormone (PTH) enhancer. Defects in CCND1 are a cause of multiple myeloma (MM) [MIM:254500]. MM is a malignant tumor of plasma cells usually arising in the bone marrow and characterized by diffuse involvement of the skeletal system, hyperglobulinemia, Bence-Jones proteinuria and anemia. |

Complications of multiple myeloma are bone pain, hypercalcemia, renal failure and spinal cord compression. The aberrant antibodies that are produced lead to impaired humoral immunity and patients have a high prevalence of infection. Amyloidosis may develop in some patients. Multiple myeloma is part of a spectrum of diseases ranging from monoclonal gammopathy of unknown significance (MGUS) to plasma cell leukemia. Note=A chromosomal aberration involving CCND1 is found in multiple myeloma. Translocation t(11;14)(q13;q32) with the IgH locus.

Sequence similarities

Belongs to the cyclin family. Cyclin D subfamily.

Post-translational modifications


Phosphorylation at Thr-286 by MAP kinases is required for ubiquitination and degradation following DNA damage. It probably plays an essential role for recognition by the FBXO31 component of SCF (SKP1-cullin-F-box) protein ligase complex.

Ubiquitinated, primarily as 'Lys-48'-linked polyubiquitination. Ubiquitinated by a SCF (SKP1-CUL1-F-box protein) ubiquitin-protein ligase complex containing FBXO4 and CRYAB (By similarity). Following DNA damage it is ubiquitinated by some SCF (SKP1-cullin-F-box) protein ligase complex containing FBXO31. Ubiquitination leads to its degradation and G1 arrest. Deubiquitinated by USP2; leading to stabilize it.

Cellular localization

Nucleus.

Anti-Cyclin D1 antibody [EPR2241] images



All lanes : Anti-Cyclin D1 antibody [EPR2241] (ab134175) at 1/10000 dilution (purified)

Lane 1 : Mouse kidney

Lane 2 : Mouse spleen

Lane 3 : Rat heart

Lysates/proteins at 20 µg per lane.

Secondary


HRP goat anti-rabbit (H+L) at 1/1000 dilution

Predicted band size : 34 kDa

Observed band size : 34 kDa


Blocking buffer: 5% NFDm/TBST

Dilution buffer: 5% NFDm/TBST




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cyclin D1 antibody [EPR2241] (ab134175)

Immunohistochemical staining of paraffin embedded human endometrial adenocarcinoma with purified ab134175 at a dilution of 1/100. A HRP goat anti-rabbit (ab97051) was used as the secondary antibody at a dilution of 1/500 and the sample was counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.



Immunocytochemistry/ Immunofluorescence - Anti-Cyclin D1 antibody [EPR2241] (ab134175)

Immunofluorescent staining of Ramos cells (fixed in 4% PFA, permeabilized with 0.1% Triton X 100) using purified ab134175 at a dilution of 1/50. An Alexa Fluor[®] 488 goat anti-rabbit antibody was used as the secondary at a dilution of 1/500 and the cells were counterstained with DAPI. The negative control is shown in the bottom right hand panel - for the negative control, Alex Fluor[®] 594 goat anti-mouse was used at a dilution of 1/500.




Immunoprecipitation - Anti-Cyclin D1 antibody [EPR2241] (ab134175)

ab134175 (purified) at 1/30 immunoprecipitating cyclin D1 in A431 cells (Lane 1). For western blotting, a HRP-conjugated goat anti-rabbit IgG, was used as the secondary antibody (1/1000).


Blocking buffer and concentration: 5% NFDN/TBST.

Diluting buffer and concentration: 5% NFDN /TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cyclin D1 antibody [EPR2241] (ab134175)


Immunohistochemical analysis of paraffin-embedded human mantle cell lymphoma tissue, labeling Cyclin D1 with unpurified ab134175 at 1/100 dilution.



Immunocytochemistry/ Immunofluorescence - Anti-Cyclin D1 antibody [EPR2241] (ab134175)

Unpurified ab134175 staining Cyclin D1 in MCF7 cells treated with KN-93 (ab120980). The cells were fixed with 100% methanol (5min) and then blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated with ab134175 at 10µg/ml and ab7291 at 1µg/ml overnight at +4°C, followed by a further incubation at room temperature for 1h with an Goat anti-Rabbit Alexa 488 secondary (ab150081) at 2 µg/ml (shown in green) and Goat anti-Mouse Alexa 594 secondary (ab150120) at 2 µg/ml (shown in pseudo color red). Nuclear DNA was labelled in blue with DAPI.

Negative controls: 1– Rabbit primary and anti-mouse secondary antibody; 2 – Mouse primary antibody and anti-rabbit secondary antibody. Controls 1 and 2 indicate that there is no unspecific reaction between primary and secondary antibodies used.



Western blot - Anti-Cyclin D1 antibody [EPR2241]
(ab134175)

All lanes : Anti-Cyclin D1 antibody
[EPR2241] (ab134175) at 1/100000 dilution
(purified)

Lane 1 : MCF-7 cell lysate

Lane 2 : LnCap cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


HRP goat anti-rabbit (H+L) at 1/1000 dilution

Predicted band size : 34 kDa

Observed band size : 34 kDa

Blocking buffer: 5% NFDm/TBST

Dilution buffer: 5% NFDm/TBST



Western blot - Anti-Cyclin D1 antibody [EPR2241]
(ab134175)

All lanes : Anti-Cyclin D1 antibody
[EPR2241] (ab134175) at 1/20000 dilution
(purified)

Lane 1 : HeLa cell lysate

Lane 2 : A431 cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


HRP goat anti-rabbit (H+L) at 1/1000 dilution

Predicted band size : 34 kDa

Observed band size : 34 kDa

Blocking buffer: 5% NFDm/TBST

Dilution buffer: 5% NFDm/TBST




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cyclin D1 antibody [EPR2241] (ab134175)

IHC image of ab134175 staining Cyclin D1 in human esophagus formalin fixed paraffin embedded tissue sections*, performed on a Leica Bond. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab134175, 5µg/ml working concentration, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with hematoxylin and mounted with DPX. No primary antibody was used in the secondary only control (shown on the inset).

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.


*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre



Immunocytochemistry/ Immunofluorescence - Anti-Cyclin D1 antibody [EPR2241] (ab134175)


This image is courtesy of an Abreview submitted by Kirk Mcmanus.

Immunocytochemical analysis of HeLa cells, labeling Cyclin D1 with ab134175 at a dilution of 1/200. Cells were paraformaldehyde fixed and permeabilized with 0.5% Triton X-100 in PBS. Incubation with the primary antibody was for 1 hour at 22°C. Cells were counterstained with DAPI following immunostaining.



IHC image of ab134175 staining Cyclin D1 in rat esophagus formalin fixed paraffin embedded tissue sections, performed on a Leica Bond. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab134175, 5µg/ml working concentration, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with hematoxylin and mounted with DPX. No primary antibody was used in the secondary only control (shown on the inset).

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Developed using the ECL technique


Performed under reducing conditions.

Predicted band size : 34 kDa

Exposure time : 3 minutes

MG-63 cells were incubated at 37°C for 24h with vehicle control (0 μ M) and different concentrations of 3-aminobenzamide (3-AB) (ab141069). Decreased expression of cyclin D1 (unpurified ab134175) in MG-63 cells correlates with an increase in 3-aminobenzamide (3-AB) concentration, as described in literature.

Whole cell lysates were prepared with RIPA buffer (containing protease inhibitors and sodium orthovanadate), 10 μ g of each were loaded on the gel and the WB was run under reducing conditions. After transfer the membrane was blocked for an hour using 5% BSA before being incubated with unpurified ab134175 at 1/500 dilution and ab8226 at 1 μ g/ml overnight at 4°C. Antibody binding was detected using an anti-rabbit HRP secondary antibody (ab97051) at 1/10000 dilution and visualised using ECL development solution.




Anti-Cyclin D1 antibody [EPR2241]
(ab134175) at 1/10000 dilution (unpurified) +
MCF7 cell lysate at 10 μ g

Secondary

HRP labelled goat anti-rabbit at 1/2000
dilution

Predicted band size : 34 kDa


Secondary goat anti-rabbit HRP (ab6721)



Immunocytochemistry/ Immunofluorescence - Anti-Cyclin D1 antibody [EPR2241] (ab134175)

This image is courtesy of an anonymous Abreview.

Immunocytochemical analysis of human MCF7 cells, labeling Cyclin D1 with ab134175 at a dilution of 1/100. Cells were paraformaldehyde fixed and permeabilized with 0.1% Triton X-100. To avoid non-specific binding, cells were blocked with 1% Goat serum + 1% BSA in PBS for 1 hour at 22°C. Incubation with the primary antibody was for 16 hours at 4°C. An Alexa Fluor® 488 conjugated goat anti-rabbit IgG polyclonal antibody, diluted 1/500.



Other - Anti-Cyclin D1 antibody [EPR2241] (ab134175)

Equilibrium dissociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors