

Product datasheet

Anti-Cytochrome P450 2B1 + 2B2 antibody [b/e3 (9.14)] ab22719

[1 Image](#)

Overview

| | |
|----------------------------|---|
| Product name | Anti-Cytochrome P450 2B1 + 2B2 antibody [b/e3 (9.14)] |
| Description | Mouse monoclonal [b/e3 (9.14)] to Cytochrome P450 2B1 + 2B2 |
| Host species | Mouse |
| Specificity | This product does not react with Rat CYP 1A1, 1A2, 2A1, 2C6, 2C7, 2C11, 4A2 and 4A3 |
| Tested applications | Suitable for: ELISA, WB, IP |
| Species reactivity | Reacts with: Mouse, Rat |
| Immunogen | Rat liver cytochrome P450 2B1 and 2B2. |

Properties

| | |
|-----------------------------|--|
| Form | Liquid |
| Storage instructions | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle. |
| Storage buffer | Preservative: None Constituents: PBS |
| Clonality | Monoclonal |
| Clone number | b/e3 (9.14) |
| Isotype | IgG1 |

Applications

Our [Abpromise guarantee](#) covers the use of **ab22719** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|-------------|-----------|--|
| ELISA | | Use at an assay dependent concentration. |
| WB | | Use at an assay dependent concentration. Predicted molecular weight: 56 kDa. |

| Application | Abreviews | Notes |
|-------------|-----------|---------------------------------|
| IP | | Use a concentration of 5 µg/ml. |

Target


Relevance

The Cytochrome P450 superfamily of enzymes is one of three enzyme systems which metabolize the fatty acid arachadonic acid (AA) to regulators of vascular tone. P450 enzymes are monooxygenase enzymes which require several co-factors such as nicotinamide adenine dinucleotide phosphate (NADPH) and P450 reductase. There are over 200 known genes which encode P450s. Epoxygenases are those P450s which metabolize AA to epoxyeicosatrienoic acid (EETs) and omega-hydroxylases are those P450s which produce 19- and 20-hydroxyeicosatetraenoic acids (19- and 20-HETE). As well as fatty acid metabolism, P450s also metabolize many drugs and toxins. Cytochrome P450 3A4 is abundantly expressed in liver and small intestine and is inducible by barbiturates, glucocorticoids and rifampicin.

Cellular localization

Membrane bound, endoplasmic reticulum

Images



Cytochrome P450 2B1 + 2B2 was immunoprecipitated using 0.5mg Mouse Liver tissue, 5µg of Mouse monoclonal to Cytochrome P450 2B1 + 2B2 and 50µl of protein G magnetic beads (+). No antibody was added to the control (-).

The antibody was incubated under agitation with Protein G beads for 10min, Mouse Liver tissue lysate diluted in RIPA buffer was added to each sample and incubated for a further 10min under agitation.

Proteins were eluted by addition of 40µl SDS loading buffer and incubated for 10min at 70°C; 10µl of each sample was separated on a SDS PAGE gel, transferred to a nitrocellulose membrane, blocked with 5% BSA and probed with ab22719.

Secondary: Goat polyclonal to mouse IgG light chain specific (HRP) at 1/20,000 dilution.

Band: 56kDa; Cytochrome P450 2B1 + 2B2

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors