

Product datasheet

Anti-Cytochrome P450 3A4 antibody ab98319

1 Image

Overview

Product name	Anti-Cytochrome P450 3A4 antibody
Description	Rabbit polyclonal to Cytochrome P450 3A4
Host species	Rabbit
Tested applications	Suitable for: WB
Species reactivity	Reacts with: Human Predicted to work with: Mouse, Rat, Sheep, Rabbit, Goat, Horse, Chicken, Guinea pig, Cow, Cat, Dog, African green monkey
Immunogen	Synthetic peptide corresponding to a region within internal amino acids 395-444 (MIPSYALHRDPKYWTEPEKFLPERFSKKNKDNIDPYYTPFGSGPRNCIG) of Human Cytochrome P450 3A4 (NP_059488). Run BLAST with ExPASy Run BLAST with NCBI
Positive control	THP-1 cell lysate.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Preservative: 0.09% Sodium azide Constituents: 2% Sucrose, PBS
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab98319** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
-------------	-----------	-------

Application	Abreviews	Notes
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: 57 kDa. Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.

Target

Function	Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It performs a variety of oxidation reactions (e.g. caffeine 8-oxidation, omeprazole sulphoxidation, midazolam 1'-hydroxylation and midazolam 4-hydroxylation) of structurally unrelated compounds, including steroids, fatty acids, and xenobiotics. The enzyme also hydroxylates etoposide.
Tissue specificity	Expressed in prostate and liver. According to some authors, it is not expressed in brain (PubMed:19094056). According to others, weak levels of expression are measured in some brain locations (PubMed:19359404 and PubMed:18545703). Also expressed in epithelium of the small intestine and large intestine, bile duct, nasal mucosa, kidney, adrenal cortex, epithelium of the gastric mucosa with intestinal metaplasia, gallbladder, intercalated ducts of the pancreas, chief cells of the parathyroid and the corpus luteum of the ovary (at protein level).
Sequence similarities	Belongs to the cytochrome P450 family.
Post-translational modifications	Polyubiquitinated in the presence of AMFR and UBE2G1 and also STUB1/CHIP and UBE2D1 (in vitro).
Cellular localization	Endoplasmic reticulum membrane. Microsome membrane.

Images



Anti-Cytochrome P450 3A4 antibody (ab98319) at 1 µg/ml (5% skim milk/PBS buffer) + THP-1 cell lysate at 10 µg

Predicted band size: 57 kDa

Gel concentration: 12%

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors