

Product datasheet

Anti-DDX4 / MVH antibody [EPR3849] ab108392

Recombinant RabMAb

1 Image

Overview

Product name	Anti-DDX4 / MVH antibody [EPR3849]
Description	Rabbit monoclonal [EPR3849] to DDX4 / MVH
Host species	Rabbit
Tested applications	Suitable for: WB, IP Unsuitable for: Flow Cyt, ICC or IHC-P
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	A synthetic peptide corresponding to residues in Human DDX4/ MVH
Positive control	Human testis, Human ovary cancer, K562, seminoma, HeLa, NCCIT and LNCaP cell lysates
General notes	

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMab[®] patents](#)

This product is a recombinant rabbit monoclonal antibody.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
Storage buffer	PBS 49%, Sodium azide 0.01%, Glycerol 50%, BSA 0.05%
Purity	Tissue culture supernatant
Clonality	Monoclonal
Clone number	EPR3849
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab108392** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/10000 - 1/50000. Detects a band of approximately 83 kDa (predicted molecular weight: 79 kDa).
IP		1/10 - 1/100.

Application notes Is unsuitable for Flow Cyt, ICC or IHC-P.

Target


Function Probable ATP-dependent RNA helicase required during spermatogenesis (PubMed:10920202, PubMed:21034600). Required to repress transposable elements and preventing their mobilization, which is essential for the germline integrity. Acts via the piRNA metabolic process, which mediates the repression of transposable elements during meiosis by forming complexes composed of piRNAs and Piwi proteins and governs the methylation and subsequent repression of transposons. Involved in the secondary piRNAs metabolic process, the production of piRNAs in fetal male germ cells through a ping-pong amplification cycle.

Tissue specificity Expressed only in ovary and testis. Expressed in migratory primordial germ cells in the region of the gonadal ridge in both sexes.

Sequence similarities Belongs to the DEAD box helicase family. DDX4/VASA subfamily.
Contains 1 helicase ATP-binding domain.
Contains 1 helicase C-terminal domain.

Cellular localization Cytoplasm. Cytoplasm, perinuclear region. Component of the meiotic nuage, also named P granule, a germ-cell-specific organelle required to repress transposon activity during meiosis.

Images



All lanes : Anti-DDX4 / MVH antibody
[EPR3849] (ab108392) at 1/10000 dilution

- Lane 1 :** Human testis lysate
- Lane 2 :** Human ovary cancer lysate
- Lane 3 :** K562 cell lysate
- Lane 4 :** Seminoma cell lysate
- Lane 5 :** HeLa cell lysate
- Lane 6 :** NCCIT cell lysate
- Lane 7 :** LNCaP cell lysate

Lysates/proteins at 10 µg per lane.

Predicted band size: 79 kDa

Observed band size: 83 kDa

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors