

Product datasheet

Anti-DHX57 antibody ab86784

[2 Images](#)

Overview

Product name	Anti-DHX57 antibody
Description	Rabbit polyclonal to DHX57
Tested applications	Suitable for: WB, IP
Species reactivity	Reacts with: Human Predicted to work with: Chimpanzee, Gorilla, Orangutan
Immunogen	Synthetic peptide corresponding to a region between residues 150 and 200 of human DHX57 (NP_945314.1).
Positive control	HeLa and 293T whole cell lysates

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Preservative: 0.09% Sodium Azide Constituents: 0.1% BSA, Tris buffered saline
Purity	Immunogen affinity purified
Purification notes	ab86784 was affinity purified using an epitope specific to DHX57 immobilized on solid support.
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab86784** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		
IP		

Application notes

IP: Use at 10 µg/mg of lysate.
WB: 1/2000 - 1/10,000. Predicted molecular weight: 156 kDa.

Not yet tested in other applications.
Optimal dilutions/concentrations should be determined by the end user.

Target


Function

Probable ATP-binding RNA helicase.

Sequence similarities

Belongs to the DEAD box helicase family. DEAH subfamily.
Contains 1 C3H1-type zinc finger.
Contains 1 helicase ATP-binding domain.
Contains 1 helicase C-terminal domain.
Contains 1 UBA domain.

Images



All lanes : Anti-DHX57 antibody (ab86784) at 0.04 µg/ml

Lane 1 : HeLa whole cell lysate at 50 µg

Lane 2 : HeLa whole cell lysate at 15 µg

Lane 3 : HeLa whole cell lysate at 5 µg


Lane 4 : 293T whole cell lysate at 50 µg

Developed using the ECL technique

Predicted band size : 156 kDa

Observed band size : 171 kDa

Exposure time : 3 minutes



Detection of DHX57 by Western Blot of Immunoprecipitate.

ab86784, at 0.4 µg/ml, staining DHX57 in HeLa whole cell lysate immunoprecipitated using ab86784 at 10µg/mg lysate (1 mg/IP; 20% of IP loaded/lane).

Detection: Chemiluminescence with an exposure time of 30 seconds.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors