

## Product datasheet

# Donkey Anti-Goat IgG H&L (DyLight® 488) preadsorbed ab96935

★★★★★ 1 Abreviews 2 Images

### Overview

<b>Product name</b>	Donkey Anti-Goat IgG H&L (DyLight® 488) preadsorbed
<b>Description</b>	Donkey polyclonal Secondary Antibody to Goat IgG - H&L (DyLight® 488), pre-adsorbed
<b>Target species</b>	Goat
<b>Specificity</b>	By immunoelectrophoresis and ELISA this antibody reacts specifically with Goat IgG and with light chains common to other Goat immunoglobulins. No antibody was detected against non immunoglobulin serum proteins. Less than 0.1% cross reactivity to chicken, horse, human, mouse, pig, rabbit and rat was detected.
<b>Tested applications</b>	<b>Suitable for:</b> Flow Cyt, IHC-P, ICC/IF
<b>Minimal cross-reactivity</b>	Chicken, Horse, Human, Mouse, Pig, Rabbit, Rat <a href="#">more details</a>
<b>Conjugation</b>	DyLight® 488. Ex: 493nm, Em: 518nm

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C.
<b>Storage buffer</b>	Preservative: 0.09% Sodium Azide Constituents: 0.2% BSA, PBS
<b>Purity</b>	Immunogen affinity purified
<b>Purification notes</b>	Antiserum was cross adsorbed using chicken, human, mouse, pig, rabbit and rat immunosorbents to remove cross reactive antibodies. This antibody was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 488.
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG


### Applications

Our [Abpromise guarantee](#) covers the use of **ab96935** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		1/50 - 1/200.


### Donkey Anti-Goat IgG H&L (DyLight® 488) preadsorbed images



Immunohistochemistry (PFA perfusion fixed frozen sections) - Donkey Anti-Goat IgG H&L (DyLight® 488) preadsorbed (ab96935)

Image is courtesy of an AbReview submitted by Ms Francoise Geffroy.

Immunohistochemical analysis of Paraformaldehyde perfusion-fixed frozen human neuroblastoma xenograft into a murine host. The image is labelling LIMP2 with [ab87637](#) at 20°C for 1 hour at a dilution of 1/50 with 5% Donkey Serum. Permeabilization was done with 0.2% Triton. Secondary used was [ab96935](#) at a 1/250 dilution.



DyLight®-Donkey polyclonal Secondary Antibody to Goat IgG - H&L (DyLight® 488), preadsorbed(ab96935)

Emission spectra of DyLight® fluorochromes available in our catalog. Line colors represent the approximate visible colors of the wavelength maxima.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

- 
- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors