

## Product datasheet

# Anti-E. coli O119 antibody ab78777

### Overview

---

<b>Product name</b>	Anti-E. coli O119 antibody
<b>Description</b>	Rabbit polyclonal to E. coli O119
<b>Specificity</b>	No cross reactivity with other serotypes observed.
<b>Tested applications</b>	<b>Suitable for:</b> Agglutination
<b>Species reactivity</b>	Reacts with E. coli O119.
<b>Immunogen</b>	Tissue/ cell preparation of E. coli O119 (heat inactivated).
<b>Positive control</b>	Antigenic suspension.
<b>General notes</b>	Reagents should be allowed to stand at 15°C to 25°C for at least 30 minutes prior to use.

### Properties

---

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	Preservative: 0.08% Sodium Azide Constituents: Whole serum
<b>Purity</b>	Whole antiserum
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG

### Applications

---

Our [Abpromise guarantee](#) covers the use of **ab78777** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Agglutination		Use at an assay dependent concentration. See attached protocol.

---

## Relevance

*Escherichia coli* is a group of gram-negative bacilli belonging to the family *Enterobacteriaceae* and is considered as one of the bacteria in the human intestinal normal microbial flora. Their serological types are determined in combination with somatic antigens (O group: O1-O173) and flagella antigen (H type: H1-H56). The *E. coli* that cause intestinal infectious diseases including diarrhea, acute gastritis or colitis are referred to as pathogenic *E. coli*. These are classified into the following four groups according to differences in the mode of pathogenicity; enteropathogenic *E. coli* (EPEC), enteroinvasive *E. coli* (EIEC), enterotoxigenic *E. coli* (ETEC) and enterohemorrhagic *E. coli* (EHEC). Although the identification of pathogenic *E. coli* requires verification of their pathogenicity, pathogenic *E. coli* often have specific serotypes; therefore, typing of the serogroup and serotype is necessary in screening pathogenic *E. coli*.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors