


Product datasheet

Anti-EYA2 antibody ab95875

3 Images

Overview

Product name	Anti-EYA2 antibody
Description	Rabbit polyclonal to EYA2
Host species	Rabbit
Tested applications	Suitable for: WB, IP, IHC-P
Species reactivity	Reacts with: Human Predicted to work with: Horse, Cow, Chimpanzee, Rhesus monkey, Gorilla, Opossum, Orangutan, Bat 
Immunogen	Synthetic peptide corresponding to a region between residue 100 and 150 of human EYA2 (NP_005235.3)
Positive control	293T cells lysate

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Preservative: 0.09% Sodium Azide Constituents: Tris citrate/phosphate, pH 7-8
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab95875** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
WB		1/2000 - 1/10000. Predicted molecular weight: 59 kDa.
IP		Use at 2-5 µg/mg of lysate.

Application	Abreviews	Notes
IHC-P		1/500 - 1/2000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Target


Function	Tyrosine phosphatase that specifically dephosphorylates 'Tyr-142' of histone H2AX (H2AXY142ph). 'Tyr-142' phosphorylation of histone H2AX plays a central role in DNA repair and acts as a mark that distinguishes between apoptotic and repair responses to genotoxic stress. Promotes efficient DNA repair by dephosphorylating H2AX, promoting the recruitment of DNA repair complexes containing MDC1. Its function as histone phosphatase probably explains its role in transcription regulation during organogenesis. Coactivates SIX1. Seems to coactivate SIX2, SIX4 and SIX5. Together with SIX1 and DACH2 seem to be involved in myogenesis. May be involved in development of the eye. Interaction with GNAZ and GNAI2 prevents nuclear translocation and transcriptional activity.
Tissue specificity	Highest expression in muscle with lower levels in kidney, placenta, pancreas, brain and heart.
Sequence similarities	Belongs to the HAD-like hydrolase superfamily. EYA family.
Developmental stage	At the begin of fourth week of development detected in cytoplasm of somite cells. Between the sixth and eighth week of development detected in cytoplasm of limb bud cells.
Cellular localization	Cytoplasm. Nucleus.

Images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast carcinoma tissue labelling EYA2 with ab95875 at 1/1000 (1µg/ml).
Detection: DAB.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EYA2 antibody (ab95875)



All lanes : Anti-EYA2 antibody (ab95875) at 0.2 µg/ml

Lane 1 : 293T cell lysate at 50 µg


Lane 2 : 293T cell lysate at 15 µg

Lane 3 : 293T cell lysate at 5 µg

Developed using the ECL technique.

Predicted band size: 59 kDa

Exposure time: 30 seconds



Detection of human Eya2 by Western Blot of

Immunoprecipitate. ab95875 at 0.1 µg/ml

staining Eya2 in 293T cell lysate

immunoprecipitated using ab95875 at 3 µg/mg

(1mg/IP; 20% of IP loaded/lane) Detection:

Chemiluminescence with exposure time of 10

seconds..

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors