

Product datasheet

Anti-Ephrin B (phospho Y317) antibody ab33473

1 Image

Overview

Product name	Anti-Ephrin B (phospho Y317) antibody
Description	Rabbit polyclonal to Ephrin B (phospho Y317)
Host species	Rabbit
Specificity	This antibody is specific to Dak Ephrin B protein phosphorylated at Tyr317.
Tested applications	Suitable for: WB
Species reactivity	Reacts with: Mouse, Rat, Chicken, Human, Xenopus laevis
Immunogen	Phosphopeptide corresponding to amino acid residues surrounding the phospho Tyr317 of Chicken EphrinB.
Positive control	Rat testes lysate.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Preservative: None Constituents: 50% Glycerol, 100µg/ml BSA, 150mM Sodium chloride, 10mM HEPES. pH 7.5
Purity	Protein A purified
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab33473** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
-------------	-----------	-------

WB

Application notes	WB: 1/1000. Detects a band of approximately 46 kDa (predicted molecular weight: 37 kDa). Note: The immunolabeling of the Ephrin B band is blocked by lambda phosphatase treatment.
--------------------------	---

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

Target


Relevance

Ephrin B proteins are thought to play key roles in cellular functions as diverse as neuronal migration and blood vessel development. Ephrin B molecules expressed at the membrane surface bind to the Ephrin B family receptors on target cells during cell to cell contact. This interaction leads to cell signaling in the target cell but also generates a reverse signal in the cell expressing Ephrin B on its surface. This reverse signaling event is thought to be critical for vessel maturation and neuronal development. Importantly, tyrosine phosphorylation of Ephrin B is thought to be a critical component of this reverse signaling event. Recent work demonstrated that Tyr331 of Ephrin B was phosphorylated in HEK293 cells after stimulation by the soluble Ephrin B2 receptor tyrosine kinase.

Cellular localization

Membrane; single pass type I membrane protein.

Images



All lanes : Anti-Ephrin B (phospho Y317) antibody (ab33473) at 1/1000 dilution

Lane 1 : Rat testes lysate treated with this Ephrin B antibody phosphorylated at Tyr317 (control).

Lane 2 : Rat testes lysate incubated in lambda phosphatase (1200 unit for 30 min) before being exposed to this Ephrin B antibody phosphorylated at Tyr317.

Lysates/proteins at 50 µg per lane.

Predicted band size: 37 kDa

Observed band size: 46 kDa

Additional bands at: 100 kDa. We are unsure as to the identity of these extra bands.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors