


Product datasheet

Anti-FADS2 antibody ab72189

1 References 2 Images

Overview

<b>Product name</b>	Anti-FADS2 antibody
<b>Description</b>	Rabbit polyclonal to FADS2
<b>Host species</b>	Rabbit
<b>Tested applications</b>	<b>Suitable for:</b> WB, IP
<b>Species reactivity</b>	<b>Reacts with:</b> Human <b>Predicted to work with:</b> Monkey, Baboon, Orangutan 
<b>Immunogen</b>	Synthetic peptide conjugated to KLH derived from within residues 1 - 100 of Human FADS2. Read Abcam's proprietary immunogen policy (Peptide available as <a href="#">ab90992</a> .)
<b>Positive control</b>	This antibody gave a positive signal in the following lysates: Human liver tissue lysate; Human brain tissue lysates; HeLa whole cell lysate; Hep G2 whole cell lysate.

Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	Preservative: 0.02% Sodium Azide Constituents: 1% BSA, PBS, pH 7.4
<b>Purity</b>	Immunogen affinity purified
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG

Applications


Our [Abpromise guarantee](#) covers the use of **ab72189** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 45 kDa (predicted molecular weight: 52 kDa).

Application	Abreviews	Notes
IP		Use at an assay dependent concentration.
<b>Target</b>		
<b>Function</b>	Component of a lipid metabolic pathway that catalyzes biosynthesis of highly unsaturated fatty acids (HUFA) from precursor essential polyunsaturated fatty acids (PUFA) linoleic acid (LA) (18:2n-6) and alpha-linolenic acid (ALA) (18:3n-3). Catalyzes the first and rate limiting step in this pathway which is the desaturation of LA (18:2n-6) and ALA (18:3n-3) into gamma-linoleic acid (GLA) (18:3n-6) and stearidonic acid (18:4n-3) respectively and other desaturation steps. Highly unsaturated fatty acids (HUFA) play pivotal roles in many biological functions. It catalyzes as well the introduction of a cis double bond in palmitate to produce the mono-unsaturated fatty acid sapienate, the most abundant fatty acid in sebum.	
<b>Tissue specificity</b>	Expressed in a wide array of tissues, highest expression is found in liver followed by brain, lung, heart, and retina. A lower level is found in breast tumor when compared with normal tissues; lowest levels were found in patients with poor prognostic index.	
<b>Pathway</b>	Lipid metabolism; polyunsaturated fatty acid biosynthesis.	
<b>Sequence similarities</b>	Belongs to the fatty acid desaturase family. Contains 1 cytochrome b5 heme-binding domain.	
<b>Developmental stage</b>	Found in fetal heart.	
<b>Domain</b>	The histidine box domains may contain the active site and/or be involved in metal ion binding.	
<b>Cellular localization</b>	Endoplasmic reticulum membrane.	

## Images



FADS2 was immunoprecipitated using 0.5mg HepG2 whole cell extract, 5µg of Rabbit polyclonal to FADS2 and 50µl of protein G magnetic beads (+). No antibody was added to the control (-).


The antibody was incubated under agitation with Protein G beads for 10min, HepG2 whole cell extract lysate diluted in RIPA buffer was added to each sample and incubated for a further 10min under agitation.

Proteins were eluted by addition of 40µl SDS loading buffer and incubated for 10min at 70°C; 10µl of each sample was separated on a SDS PAGE gel, transferred to a nitrocellulose membrane, blocked with 5% BSA and probed with ab72189.

Secondary: Mouse monoclonal [SB62a]

Secondary Antibody to Rabbit IgG light chain (HRP) ([ab99697](#)).

Band: 45kDa:FADS2.



Western blot - FADS2 antibody (ab72189)

**All lanes** : Anti-FADS2 antibody (ab72189) at 1 µg/ml

**Lane 1** : HeLa (Human epithelial carcinoma cell line) Whole Cell Lysate

**Lane 2** : Human liver tissue lysate - total protein ([ab29889](#))

**Lane 3** : HepG2 (Human hepatocellular liver carcinoma cell line) Whole Cell Lysate

**Lane 4** : Human brain tissue lysate - total protein ([ab29466](#))

Lysates/proteins at 10 µg per lane.

### Secondary

**All lanes** : Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

**Predicted band size:** 52 kDa

**Observed band size:** 45 kDa

**Additional bands at:** 62.5 kDa. We are unsure as to the identity of these extra bands.

**Exposure time:** 2 minutes

ab72189 detects a strong band at 45 kDa in human liver and brain tissue lysates, as well as HeLa and HepG2 whole cell lysates. This antibody was raised against an immunogen that is predicted to cross react with all known isoforms of Fatty acid desaturase 2 (FADS2). FADS2 has three isoforms, isoform one has a molecular weight of 52 kDa with isoforms two and three a molecular weight of 49 and 45 respectively (SwissProt data). The band we have observed is thought to represent isoform 3 of FADS2 (45 kDa).

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors