

Product datasheet

Anti-GRK1 antibody [D11] ab2776

[3 References](#) [4 Images](#)

Overview

Product name	Anti-GRK1 antibody [D11]
Description	Mouse monoclonal [D11] to GRK1
Host species	Mouse
Specificity	Detects G-protein-associated rhodopsin kinase 1a/b (GRK1a/b) from human and bovine tissues. This antibody does not cross-react with other G protein-coupled kinases.
Tested applications	Suitable for: ICC/IF, IHC-Fr, WB, IP
Species reactivity	Reacts with: Mouse, Cow, Human
Immunogen	Full length native protein (purified) corresponding to Human GRK1.
Epitope	This antibody recognizes an N-terminal epitope of human GRK1 which is conserved in both GRK1a and GRK1b.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.05% Sodium azide
Purity	Protein G purified
Primary antibody notes	The sensation of sight is the result of a cascade of events starting with the interaction of photoactivated rhodopsin with a protein called transducin. Rhodopsin kinase is a G-protein-coupled Ser/Thr kinase, also known as GRK1, which is a key element in the regulation of this cascade. Following phosphorylation by GRK1, arrestin is recruited to phospho-rhodopsin quenching its phototransductive activity by preventing further interaction with transducin. By breaking the cycle of phototransduction, GRK1 plays an important role in the restoration of the system for subsequent visual events.
Clonality	Monoclonal
Clone number	D11
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab2776** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		1/20 - 1/200.
IHC-Fr		1/1000.
WB		1/1000.
IP		Use at an assay dependent concentration.

Target


Function	Retina-specific kinase involved in the signal turnoff via phosphorylation of rhodopsin (RHO), the G protein- coupled receptor that initiates the phototransduction cascade. This rapid desensitization is essential for scotopic vision and permits rapid adaptation to changes in illumination.
Tissue specificity	Retinal-specific. Expressed in rods and cones cells.
Involvement in disease	Night blindness, congenital stationary, Oguchi type 2
Sequence similarities	Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. GPRK subfamily. Contains 1 AGC-kinase C-terminal domain. Contains 1 protein kinase domain. Contains 1 RGS domain.
Post-translational modifications	Autophosphorylated, Ser-21 is a minor site of autophosphorylation compared to Ser-491 and Thr-492 (By similarity). Phosphorylation at Ser-21 is regulated by light and activated by cAMP. Farnesylation is required for full activity.
Cellular localization	Membrane.

Images




Immunocytochemistry/ Immunofluorescence - Anti-GRK1 antibody [D11] (ab2776)

Immunocytochemistry/Immunofluorescence analysis of GRK1 (green) in murine cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes at room temperature and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were incubated with ab2776 (1:200) overnight in a humidified chamber. Cells were washed with PBST and incubated with a DyLight-conjugated secondary antibody for 45 minutes at room temperature in the dark. F-actin (red) was stained with a fluorescent phalloidin and nuclei (blue) were stained with DAPI. Images were taken at a 60X magnification.




Immunocytochemistry/ Immunofluorescence - Anti-GRK1 antibody [D11] (ab2776)

Immunocytochemistry/Immunofluorescence analysis of GRK1 (green) in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes at room temperature and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were incubated with ab2776 (1:200) overnight in a humidified chamber. Cells were washed with PBST and incubated with a DyLight-conjugated secondary antibody for 45 minutes at room temperature in the dark. F-actin (red) was stained with a fluorescent phalloidin and nuclei (blue) were stained with DAPI. Images were taken at a 60X magnification.



Immunocytochemistry/ Immunofluorescence - Anti-GRK1 antibody [D11] (ab2776)

Immunocytochemistry/Immunofluorescence analysis of GRK1 (green) in human cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes at room temperature and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were incubated with ab2776 (1:20) overnight in a humidified chamber. Cells were washed with PBST and incubated with a DyLight-conjugated secondary antibody for 45 minutes at room temperature in the dark. F-actin (red) was stained with a fluorescent phalloidin and nuclei (blue) were stained with DAPI. Images were taken at a 60X magnification.



Immunocytochemistry/ Immunofluorescence - Anti-GRK1 antibody [D11] (ab2776)

Immunolocalization of GRK1 in bovine retina using ab2776.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors