

## Product datasheet

# Anti-HLA-DR antibody [GRB-1] (PerCP) ab91333

★★★★★ 1 Abreviews 1 Image

### Overview

<b>Product name</b>	Anti-HLA-DR antibody [GRB-1] (PerCP)
<b>Description</b>	Mouse monoclonal [GRB-1] to HLA-DR (PerCP)
<b>Conjugation</b>	PerCP. Ex: 482nm, Em: 675nm
<b>Specificity</b>	Reacts with cells of the monocytic lineage, with myeloblasts and promyelocytes and cells of the B lymphocyte lineage. Does not react with polymorphonuclear leukocytes and platelets.
<b>Tested applications</b>	<b>Suitable for:</b> Flow Cyt
<b>Species reactivity</b>	<b>Reacts with:</b> Human
<b>Immunogen</b>	Tissue, cells or virus corresponding to Human HLA-DR.
<b>Positive control</b>	Normal EDTA-anticoagulated peripheral blood. Nalm-6 cell line.

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C.
<b>Storage buffer</b>	Preservative: 0.09% Sodium Azide Constituents: 1% BSA, pH 7.2
<b>Purity</b>	Immunogen affinity purified
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	GRB-1
<b>Isotype</b>	IgG2a

### Applications

Our [Abpromise guarantee](#) covers the use of **ab91333** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt	★★★★★	Use at an assay dependent concentration. <a href="#">ab110422</a> -Mouse monoclonal IgG2a, is suitable for use as an isotype control with this antibody.

## Target

---

### Function

Binds peptides derived from antigens that access the endocytic route of antigen presenting cells (APC) and presents them on the cell surface for recognition by the CD4 T-cells. The peptide binding cleft accommodates peptides of 10-30 residues. The peptides presented by MHC class II molecules are generated mostly by degradation of proteins that access the endocytic route, where they are processed by lysosomal proteases and other hydrolases. Exogenous antigens that have been endocytosed by the APC are thus readily available for presentation via MHC II molecules, and for this reason this antigen presentation pathway is usually referred to as exogenous. As membrane proteins on their way to degradation in lysosomes as part of their normal turn-over are also contained in the endosomal/lysosomal compartments, exogenous antigens must compete with those derived from endogenous components. Autophagy is also a source of endogenous peptides, autophagosomes constitutively fuse with MHC class II loading compartments. In addition to APCs, other cells of the gastrointestinal tract, such as epithelial cells, express MHC class II molecules and CD74 and act as APCs, which is an unusual trait of the GI tract. To produce a MHC class II molecule that presents an antigen, three MHC class II molecules (heterodimers of an alpha and a beta chain) associate with a CD74 trimer in the ER to form a heterononamer. Soon after the entry of this complex into the endosomal/lysosomal system where antigen processing occurs, CD74 undergoes a sequential degradation by various proteases, including CTSS and CTSL, leaving a small fragment termed CLIP (class-II-associated invariant chain peptide). The removal of CLIP is facilitated by HLA-DM via direct binding to the alpha-beta-CLIP complex so that CLIP is released. HLA-DM stabilizes MHC class II molecules until primary high affinity antigenic peptides are bound. The MHC II molecule bound to a peptide is then transported to the cell membrane surface. In B-cells, the interaction between HLA-DM and MHC class II molecules is regulated by HLA-DO. Primary dendritic cells (DCs) also to express HLA-DO. Lysosomal microenvironment has been implicated in the regulation of antigen loading into MHC II molecules, increased acidification produces increased proteolysis and efficient peptide loading.

### Sequence similarities

Belongs to the MHC class II family.  
Contains 1 Ig-like C1-type (immunoglobulin-like) domain.

### Post-translational modifications


Ubiquitinated by MARCH1 or MARCH8 at Lys-244 leading to down-regulation of MHC class II. When associated with ubiquitination of the beta subunit of HLA-DR: HLA-DRB4 'Lys-254', the down-regulation of MHC class II may be highly effective.

### Cellular localization

Cell membrane. Endoplasmic reticulum membrane. Golgi apparatus > trans-Golgi network membrane. Endosome membrane. Lysosome membrane. Late endosome membrane. The MHC class II complex transits through a number of intracellular compartments in the endocytic pathway until it reaches the cell membrane for antigen presentation.

---

## Images



Flow Cytometry - Anti-HLA-DR antibody [GRB-1]  
(PerCP) (ab91333)

This image is courtesy of an anonymous Abreview

Flow cytometry analysis of Human peripheral blood cells, staining HLA DR with ab91333.

PBMCs were isolated using a Ficoll gradient and fixed using paraformaldehyde. The sample was incubated with the primary antibody (1/100 in 2% human serum, 0.5 mM EDTA in PBS) for 30 minutes at 4°C.

**Gating Strategy:** FSC/SSC lymphocytes and then FSC-A/FSC-H singlets

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors