

Product datasheet

Anti-Hsp60 antibody [Mab11-13] ab13532

10 References 2 Images

Overview

Product name	Anti-Hsp60 antibody [Mab11-13]
Description	Mouse monoclonal [Mab11-13] to Hsp60
Host species	Mouse
Tested applications	Suitable for: WB, IP, ICC, Flow Cyt, Electron Microscopy, IHC-Fr, ELISA
Species reactivity	Reacts with: Mouse, Rat, Rabbit, Guinea pig, Hamster, Cow, Dog, Human, Pig, Drosophila melanogaster, Monkey, Snake, Dolphin, Rainbow trout Does not react with: Saccharomyces cerevisiae, Escherichia coli
Immunogen	Recombinant fragment, amino acids 31-547 (Human).
Epitope	This antibody recognizes a surface epitope of Hsp60 in the region of amino acids 288-366.
Positive control	HeLa Heat Shocked Cell Lysate or Recombinant Human Hsp60 Protein IF/ICC: HeLa cell line.
General notes	Abcam is committed to meeting high standards of ethical manufacturing and as such, we will be discontinuing this product, which has been generated by the ascites method, within the next year. We are sorry for any inconvenience this may cause. If you would like help finding an alternative product, please do not hesitate to contact our scientific support team.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.09% Sodium Azide Constituents: 50% Glycerol, PBS, pH 7.2
Purity	Protein G purified
Purification notes	This antibody has been affinity purified.
Clonality	Monoclonal
Clone number	Mab11-13
Isotype	IgG2a

Applications

Our [Abpromise guarantee](#) covers the use of **ab13532** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/10000. Detects a band of approximately 60 kDa (predicted molecular weight: 68.8 kDa).
IP		Use at an assay dependent concentration.
ICC		Use at an assay dependent concentration.
Flow Cyt		Use 0.5µg for 10 ⁶ cells. ab170191 - Mouse monoclonal IgG2a, is suitable for use as an isotype control with this antibody.
Electron Microscopy		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration.
ELISA		Use at an assay dependent concentration.

Target


Function	Implicated in mitochondrial protein import and macromolecular assembly. May facilitate the correct folding of imported proteins. May also prevent misfolding and promote the refolding and proper assembly of unfolded polypeptides generated under stress conditions in the mitochondrial matrix.
Involvement in disease	Defects in HSPD1 are a cause of spastic paraplegia autosomal dominant type 13 (SPG13) [MIM:605280]. Spastic paraplegia is a degenerative spinal cord disorder characterized by a slow, gradual, progressive weakness and spasticity of the lower limbs. Defects in HSPD1 are the cause of leukodystrophy hypomyelinating type 4 (HLD4) [MIM:612233]; also called mitochondrial HSP60 chaperonopathy or MitCHAP-60 disease. HLD4 is a severe autosomal recessive hypomyelinating leukodystrophy. Clinically characterized by infantile-onset rotary nystagmus, progressive spastic paraplegia, neurologic regression, motor impairment, profound mental retardation. Death usually occurs within the first two decades of life.
Sequence similarities	Belongs to the chaperonin (HSP60) family.
Cellular localization	Mitochondrion matrix.

Images



Immunocytochemistry/ Immunofluorescence - Anti-Hsp60 antibody [Mab11-13] (ab13532)

ICC/IF image of ab13532 stained HeLa cells. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab13532, 5µg/ml) overnight at +4°C. The secondary antibody (green) was ab96879, DyLight® 488 goat anti-mouse IgG(H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Flow Cytometry - Anti-Hsp60 antibody [Mab11-13] (ab13532)

Overlay histogram showing HeLa cells stained with ab13532 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab13532, 0.5µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) (ab96879) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2a [ICIGG2A] (ab91361, 1µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors