

## Product datasheet

# Human HIF3 alpha peptide ab20077

### Overview

---

**Product name** Human HIF3 alpha peptide

### Description

---

**Nature** Synthetic

### Amino Acid Sequence

**Species** Human

### Specifications

---

Our [Abpromise guarantee](#) covers the use of **ab20077** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

**Applications** Blocking - Blocking peptide for Anti-HIF3 alpha antibody ([ab10134](#))

**Form** Liquid

### Additional notes

- *First try to dissolve a small amount of peptide in either water or buffer. The more charged residues on a peptide, the more soluble it is in aqueous solutions.*
- *If the peptide doesn't dissolve try an organic solvent e.g. DMSO, then dilute using water or buffer.*
- *Consider that any solvent used must be compatible with your assay. If a peptide does not dissolve and you need to recover it, lyophilise to remove the solvent.*
- *Gentle warming and sonication can effectively aid peptide solubilisation. If the solution is cloudy or has gelled the peptide may be in suspension rather than solubilised.*
- *Peptides containing cysteine are easily oxidised, so should be prepared in solution just prior to use.*

### Preparation and Storage

---

**Stability and Storage** Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

Information available upon request.

### General Info

<b>Function</b>	Involves in adaptive response to hypoxia. Suppresses hypoxia-inducible expression of HIF1A and EPAS1. Binds to core DNA sequence 5'-TACGTG-3' within the hypoxia response element (HRE) of target gene promoters. The complex HIF3A-ARNT activates the transcription of reporter genes driven by HRE. Isoform 4 has a dominant-negative function of inactivating HIF1A-mediated transcription. Isoform 4 attenuates the binding of HIF1A to hypoxia-responsive elements (HRE), thus inhibiting HRE-driven transcription. Hypoxia induces down-regulation of isoform 4, leading to activation of HIF1A in hypoxia. Conversely, upon restoring normoxia, the expression of isoform 4 increases and thereby secure an inhibition of HIF1A activity. Isoform 4 may be a negative regulator of hypoxia-inducible gene expression in the kidney and may be involved in renal tumorigenesis. Functions as an inhibitor of angiogenesis in the cornea.
<b>Tissue specificity</b>	Expressed in kidney. Expressed abundantly in lung epithelial cells. Expression is regulated in an oxygen-dependent manner.
<b>Sequence similarities</b>	Contains 1 basic helix-loop-helix (bHLH) domain. Contains 2 PAS (PER-ARNT-SIM) domains.
<b>Post-translational modifications</b>	In normoxia, hydroxylated on Pro-492 in the oxygen-dependent degradation domain (ODD) by PHD. The hydroxylated proline promotes interaction with VHL, initiating rapid ubiquitination and subsequent proteasomal degradation.
<b>Cellular localization</b>	Nucleus. Cytoplasm. In the nuclei of all periportal and perivenous hepatocytes. In the distal perivenous zone, detected in the cytoplasm of the hepatocytes.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors