

## Product datasheet

# Human PUF60 peptide **ab45483**

### Overview

---

**Product name** Human PUF60 peptide

### Description

---

**Nature** Synthetic

#### Amino Acid Sequence

**Species** Human

**Sequence** YDQERFDNSDLA

### Specifications

---

Our [Abpromise guarantee](#) covers the use of **ab45483** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

**Applications** Blocking - Blocking peptide for Anti-PUF60 antibody ([ab22819](#))

**Form** Liquid

### Preparation and Storage

---

**Stability and Storage** Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

### General Info

---

**Function** DNA- and RNA-binding protein, involved in several nuclear processes such as pre-mRNA splicing, apoptosis and transcription regulation. In association with FUBP1 regulates MYC transcription at the P2 promoter through the core-TFIID basal transcription factor. Acts as a transcriptional repressor through the core-TFIID basal transcription factor. Represses FUBP1-induced transcriptional activation but not basal transcription. Decreases ERCC3 helicase activity. Does not repress TFIID-mediated transcription in xeroderma pigmentosum complementation group B (XPB) cells. Is also involved in pre-mRNA splicing. Promotes splicing of an intron with weak 3'-splice site and pyrimidine tract in a cooperative manner with U2AF2. Involved in apoptosis induction when overexpressed in HeLa cells. Isoform 6 failed to repress MYC transcription and inhibited FIR-induced apoptosis in colorectal cancer. Isoform 6 may contribute to tumor progression by enabling increased MYC expression and greater resistance

to apoptosis in tumors than in normal cells. Modulates alternative splicing of several mRNAs. Binds to relaxed DNA of active promoter regions. Binds to the pyrimidine tract and 3'-splice site regions of pre-mRNA; binding is enhanced in presence of U2AF2. Binds to Y5 RNA in association with TROVE2. Binds to poly(U) RNA.

**Tissue specificity**

Isoform 2 is expressed in colonic epithelium and colorectal epithelium cancer (at protein level). Isoform 6 is expressed in colorectal epithelial cancer but below detection level in colonic epithelium. Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine, colon and peripheral blood leukocytes.

**Sequence similarities**

Belongs to the RRM half pint family.  
Contains 3 RRM (RNA recognition motif) domains.

**Domain**

The third RNA recognition motif, called PUMP domain, is atypical and may rather mediate homodimerization and/or protein-protein interactions.

**Cellular localization**

Nucleus. Colocalizes partially with TROVE2.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

**Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

**Terms and conditions**

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors