

Product datasheet

Anti-Leptin Receptor antibody ab7212

1 References

Overview

Product name	Anti-Leptin Receptor antibody
Description	Rabbit polyclonal to Leptin Receptor
Host species	Rabbit
Specificity	Recombinant and natural mouse OB receptor.
Tested applications	Suitable for: WB
Species reactivity	Reacts with: Mouse
Immunogen	Recombinant fragment, corresponding to amino acids 217-376 of Mouse Leptin Receptor. Expressed in <i>E. coli</i>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.1% Sodium Azide Constituents: PBS
Purity	Protein A purified
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab7212** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
-------------	-----------	-------

WB

Application notes

WB: 1/2000. Predicted molecular weight: 120-140 kDa. The native leptin receptor is a homodimer held together with disulfide bonds. On a reducing SDS-PAGE, bands may appear as approximately 60 kDa subunits. Further variations in size is possible and is due to

glycosylation.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

Target

Function	Receptor for obesity factor (leptin). On ligand binding, mediates signaling through JAK2/STAT3. Involved in the regulation of fat metabolism and, in a hematopoietic pathway, required for normal lymphopoiesis. May play a role in reproduction. Can also mediate the ERK/FOS signaling pathway.
Tissue specificity	Isoform A is expressed in fetal liver and in hematopoietic tissues and choroid plexus. In adults highest expression in heart, liver, small intestine, prostate and ovary. Low level in lung and kidney. Isoform B is highly expressed in hypothalamus.
Sequence similarities	Belongs to the type I cytokine receptor family. Type 2 subfamily. Contains 4 fibronectin type-III domains. Contains 1 Ig-like (immunoglobulin-like) domain.
Domain	The cytoplasmic domain may be essential for intracellular signal transduction by activation of JAK tyrosine kinase and STATs. The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding. The box 1 motif is required for JAK interaction and/or activation.
Post-translational modifications	On ligand binding, phosphorylated on two conserved C-terminal tyrosine residues (isoform B only) by JAK2. Tyr-986 is required for complete binding and activation of PTPN11, ERK/FOS activation and, for interaction with SOCS3 (By similarity). Phosphorylation on Tyr-1141 is required for STAT3 binding/activation.
Cellular localization	Secreted and Cell membrane.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors