

Product datasheet

Anti-MCP2 antibody (Biotin) ab39627

Overview

<b>Product name</b>	Anti-MCP2 antibody (Biotin)
<b>Description</b>	Rabbit polyclonal to MCP2 (Biotin)
<b>Host species</b>	Rabbit
<b>Conjugation</b>	Biotin
<b>Tested applications</b>	<b>Suitable for:</b> Sandwich ELISA, WB
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse
<b>Immunogen</b>	Recombinant full length protein (Mouse)

Properties

<b>Form</b>	Lyophilised:Reconstitute to a concentration of 50µg/ml with sterile PBS solution containing 0.1% BSA
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	Preservative: None Constituents: 0.1% BSA, PBS
<b>Purity</b>	Immunogen affinity purified
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab39627** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Sandwich ELISA		
WB		

<b>Application notes</b>	ELISA(direct): Use at a concentration of 0.15 - 0.3 µg/ml. WB: Use at a concentration of 0.1 - 0.2 µg/ml. Predicted molecular weight: 11 kDa.
--------------------------	--

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

---

## Target

<b>Function</b>	Chemotactic factor that attracts monocytes, lymphocytes, basophils and eosinophils. May play a role in neoplasia and inflammatory host responses. This protein can bind heparin. The processed form MCP-2(6-76) does not show monocyte chemotactic activity, but inhibits the chemotactic effect most predominantly of CCL7, and also of CCL2 and CCL5 and CCL8.
<b>Tissue specificity</b>	Highest expression found in the small intestine and peripheral blood cells. Intermediate levels seen in the heart, placenta, lung, skeletal muscle, thymus, colon, ovary, spinal cord and pancreas. Low levels seen in the brain, liver, spleen and prostate.
<b>Sequence similarities</b>	Belongs to the intercrine beta (chemokine CC) family.
<b>Post-translational modifications</b>	N-terminal processed form MCP-2(6-76) is produced by proteolytic cleavage after secretion from peripheral blood monocytes.
<b>Cellular localization</b>	Secreted.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

---

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

---

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors