

Product datasheet

Anti-MVP antibody [LMR5] ab3374

1 References

Overview

Product name	Anti-MVP antibody [LMR5]
Description	Rat monoclonal [LMR5] to MVP
Host species	Rat
Tested applications	Suitable for: Flow Cyt, IHC-P, IHC-Fr
Species reactivity	Reacts with: Human
Immunogen	Tissue/ cell preparation; 2R120 cell line, derived by stepwise Doxorubicin exposure of the SW-1573 non-small cell lung carcinoma cell line.
Positive control	Positive tissue - Cell Lines: Drug-sensitive parental cell lines and their multidrugresistant derivatives. Negative control antibody: Normal rat serum.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	Constituents: 0.3% Tris HCl, 15% Sodium chloride
Purity	Tissue culture supernatant
Clonality	Monoclonal
Clone number	LMR5
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab3374** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
-------------	-----------	-------

Application	Abreviews	Notes
Flow Cyt		1/20 - 1/30. Cell permeabilization required. ab37371 - Rat polyclonal IgG, is suitable for use as an isotype control with this antibody.
IHC-P		1/20 - 1/30.
IHC-Fr		1/20 - 1/30.

Target

Function	Required for normal vault structure. Vaults are multi-subunit structures that may act as scaffolds for proteins involved in signal transduction. Vaults may also play a role in nucleo-cytoplasmic transport. Down-regulates INFG-mediated STAT1 signaling and subsequent activation of JAK. Down-regulates SRC activity and signaling through MAP kinases.
Tissue specificity	Present in most normal tissues. Higher expression observed in epithelial cells with secretory and excretory functions, as well as in cells chronically exposed to xenobiotics, such as bronchial cells and cells lining the intestine. Overexpressed in many multidrug-resistant cancer cells.
Sequence similarities	Contains 9 MVP (vault) repeats.
Domain	MVP 3 mediates interaction with PTEN. MVP 4 mediates interaction with PARP4.
Post-translational modifications	Phosphorylated on Tyr residues after EGF stimulation. Dephosphorylated by PTPN11.
Cellular localization	Cytoplasm. Nucleus > nuclear pore complex. 5% found in the nuclear pore complex. Translocates from the nucleus to the cytoplasm upon EGF treatment.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors