

Product datasheet

Anti-NM23A antibody [EPR3036] ab92327

Recombinant RabMAB

3 Images

Overview

Product name	Anti-NM23A antibody [EPR3036]
Description	Rabbit monoclonal [EPR3036] to NM23A
Host species	Rabbit
Tested applications	Suitable for: WB, IHC-P, Flow Cyt, ICC/IF Unsuitable for: IP
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide corresponding to residues in Human NM23A.
Positive control	WB: MCF7, A375, HeLa and Jurkat cell lysates IHC-P: Human lung adenocarcinoma tissue
General notes	This product is a recombinant rabbit monoclonal antibody. Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information. Our RabMAB [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMab[®] patents

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	PBS 49%,Sodium azide 0.01%,Glycerol 50%,BSA 0.05%
Purity	Tissue culture supernatant
Clonality	Monoclonal
Clone number	EPR3036
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab92327** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/1000 - 1/5000. Predicted molecular weight: 17 kDa.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol. Overnight incubation is recommended.
Flow Cyt		Use at an assay dependent concentration.
ICC/IF		1/100.

Application notes Is unsuitable for IP.

Target


Function Major role in the synthesis of nucleoside triphosphates other than ATP. Possesses nucleoside-diphosphate kinase, serine/threonine-specific protein kinase, geranyl and farnesyl pyrophosphate kinase, histidine protein kinase and 3'-5' exonuclease activities. Involved in cell proliferation, differentiation and development, signal transduction, G protein-coupled receptor endocytosis, and gene expression. Required for neural development including neural patterning and cell fate determination.

Tissue specificity Isoform 1 is expressed in heart, brain, placenta, lung, liver, skeletal muscle, pancreas, spleen and thymus. Expressed in lung carcinoma cell lines but not in normal lung tissues. Isoform 2 is ubiquitously expressed and its expression is also related to tumor differentiation. Isoform 3 is ubiquitously expressed.

Sequence similarities Belongs to the NDK family.


Cellular localization Cytoplasm. Nucleus. Cell-cycle dependent nuclear localization which can be induced by interaction with Epstein-barr viral proteins or by degradation of the SET complex by GzmA.

Images



Flow Cytometry - Anti-NM23A antibody [EPR3036]
(ab92327)

Flow Cytometry analysis of HeLa (human cervix adenocarcinoma) cells labeling NM23A with unpurified ab92327 at 1/50 dilution (10ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti-rabbit IgG (Alexa Fluor® 488)(1/2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control, cells without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.



Western blot - NM23A antibody [EPR3036]
(ab92327)

All lanes : Anti-NM23A antibody [EPR3036]
(ab92327) at 1/5000 dilution

Lane 1 : MCF-7 cell lysate

Lane 2 : A375 cell lysate

Lane 3 : HeLa cell lysate


Lane 4 : Jurkat cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP labelled goat anti-rabbit at
1/2000 dilution

Predicted band size: 17 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - NM23A antibody [EPR3036]
(ab92327)

ab92327, at a dilution of 1/100, staining
NM23A in formalin fixed, paraffin embedded
Human lung adenocarcinoma tissue by
Immunohistochemistry. Detection: DAB
staining

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors