

Product datasheet

Anti-NMNAT3 antibody ab121030

1 Image

Overview

Product name	Anti-NMNAT3 antibody
Description	Goat polyclonal to NMNAT3
Tested applications	Suitable for: WB
Species reactivity	Reacts with: Mouse Predicted to work with: Rat
Immunogen	Synthetic peptide: C-KALGQGQSVKYLLE , corresponding to internal sequence amino acids 206-220 of Mouse NMNAT3 (NP_653116.1). Run BLAST with Run BLAST with
Positive control	Mouse heart lysate

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 99% Tris buffered saline, 0.5% BSA
Purity	Immunogen affinity purified
Purification notes	Purified from goat serum by ammonium sulphate precipitation, followed by antigen affinity chromatography using the immunizing peptide.
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab121030** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use a concentration of 0.5 - 2 µg/ml. Detects a band of approximately 28 kDa (predicted molecular weight: 28 kDa). Can be blocked with ab185635 .

Target

Function	Catalyzes the formation of NAD(+) from nicotinamide mononucleotide (NMN) and ATP. Can also use the deamidated form; nicotinic acid mononucleotide (NaMN) as substrate with the same efficiency. Can use triazofurin monophosphate (TrMP) as substrate. Can also use GTP and ITP as nucleotide donors. Also catalyzes the reverse reaction, i.e. the pyrophosphorolytic cleavage of NAD(+). For the pyrophosphorolytic activity, can use NAD (+), NADH, NAAD, nicotinic acid adenine dinucleotide phosphate (NAD), nicotinamide guanine dinucleotide (NGD) as substrates. Fails to cleave phosphorylated dinucleotides NADP(+), NADPH and NAADP(+). Protects against axonal degeneration following injury.
Tissue specificity	Expressed in lung and spleen with lower levels in placenta and kidney.
Pathway	Cofactor biosynthesis; NAD(+) biosynthesis; NAD(+) from nicotinamide D-ribonucleotide: step 1/1.
Sequence similarities	Belongs to the eukaryotic NMN adenylyltransferase family.
Cellular localization	Mitochondrion.

Images



Anti-NMNAT3 antibody (ab121030) at 0.5 µg/ml + Mouse heart lysate at 35 µg
Developed using the ECL technique

Predicted band size : 28 kDa

Observed band size : 28 kDa

Additional bands at : 75 kDa. We are unsure as to the identity of these extra bands.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors