

## Product datasheet

# Natural mouse Factor Xa protein ab95261

### Overview

---

<b>Product name</b>	Natural mouse Factor Xa protein
<b>Protein length</b>	Full length protein

### Description

---

<b>Nature</b>	Native
<b>Source</b>	Native

### Amino Acid Sequence

<b>Species</b>	Mouse
----------------	-------

### Specifications

---

Our [Abpromise guarantee](#) covers the use of **ab95261** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

**Biological activity** Activity is measured in a Factor Xa clotting assay and/or chromogenic substrate assay.

**Applications** Functional Studies  
SDS-PAGE

**Purity** > 95 % SDS-PAGE.  
ab95261 is prepared by activating purified factor X with the factor X activator isolated from Russell's viper venom. Factor Xa is purified from the activation mixture by chromatography over benzamidine-Sepharose. Purity is determined by SDS-PAGE analysis.

**Form** Liquid

### Preparation and Storage

---

**Stability and Storage** Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.

Preservative: None  
Constituents: 50% Glycerol

This product is an active protein and may elicit a biological response in vivo, handle with caution.

### General Info

---

<b>Function</b>	Factor Xa is a vitamin K-dependent glycoprotein that converts prothrombin to thrombin in the presence of factor Va, calcium and phospholipid during blood clotting.
<b>Tissue specificity</b>	Plasma; synthesized in the liver.
<b>Sequence similarities</b>	Belongs to the peptidase S1 family. Contains 2 EGF-like domains. Contains 1 Gla (gamma-carboxy-glutamate) domain. Contains 1 peptidase S1 domain.
<b>Post-translational modifications</b>	The vitamin K-dependent, enzymatic carboxylation of some glutamate residues allows the modified protein to bind calcium. N- and O-glycosylated. The activation peptide is cleaved by factor IXa (in the intrinsic pathway), or by factor VIIa (in the extrinsic pathway). The iron and 2-oxoglutarate dependent 3-hydroxylation of aspartate and asparagine is (R) stereospecific within EGF domains.
<b>Cellular localization</b>	Secreted.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors