

Product datasheet

Anti-PICP antibody [8] ab17202

Overview

Product name	Anti-PICP antibody [8]
Description	Mouse monoclonal [8] to PICP
Host species	Mouse
Specificity	ab17202 is specific for native PICP from amniotic fluid.
Tested applications	Suitable for: IP, ELISA, Sandwich ELISA
Species reactivity	Reacts with: Human
Immunogen	PICP purified from second trimester amniotic fluid by affinity chromatography.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Preservative: 15mM Sodium Azide Constituents: 0.5M Sodium chloride, 0.01M PBS, pH 7.4
Purity	Protein G purified
Clonality	Monoclonal
Clone number	8
Isotype	IgG1
Light chain type	kappa

Applications

Our [Abpromise guarantee](#) covers the use of **ab17202** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IP		Use at an assay dependent concentration.
ELISA		1/6000.

Application	Abreviews	Notes
Sandwich ELISA		Use at an assay dependent concentration. Can be paired for Sandwich ELISA with Mouse monoclonal [002-18] to PICP (ab76102) . Use as Capture antibody with recommended pair.

Target

Function	Type I collagen is a member of group I collagen (fibrillar forming collagen).
Tissue specificity	Forms the fibrils of tendon, ligaments and bones. In bones the fibrils are mineralized with calcium hydroxyapatite.
Involvement in disease	Caffey disease Ehlers-Danlos syndrome 1 Ehlers-Danlos syndrome 7A Osteogenesis imperfecta 1 Osteogenesis imperfecta 2 Osteogenesis imperfecta 3 Osteogenesis imperfecta 4 Osteoporosis A chromosomal aberration involving COL1A1 is found in dermatofibrosarcoma protuberans. Translocation t(17;22)(q22;q13) with PDGF.
Sequence similarities	Belongs to the fibrillar collagen family. Contains 1 fibrillar collagen NC1 domain. Contains 1 VWFC domain.
Domain	The C-terminal propeptide, also known as COLFI domain, have crucial roles in tissue growth and repair by controlling both the intracellular assembly of procollagen molecules and the extracellular assembly of collagen fibrils. It binds a calcium ion which is essential for its function.
Post-translational modifications	Proline residues at the third position of the tripeptide repeating unit (G-X-P) are hydroxylated in some or all of the chains. Proline residues at the second position of the tripeptide repeating unit (G-P-X) are hydroxylated in some of the chains. O-linked glycan consists of a Glc-Gal disaccharide bound to the oxygen atom of a post-translationally added hydroxyl group.
Cellular localization	Secreted > extracellular space > extracellular matrix.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors