

Product datasheet

Recombinant Cat Interferon gamma protein ab93910

Overview

Product name	Recombinant Cat Interferon gamma protein
Protein length	Full length protein

Description

Nature Recombinant

Source Yeast

Amino Acid Sequence

Species Cat

Sequence

```

QAMFFKEIEE LKGYFNASNP DVADGGSLFV
DILKNWKEES DKTIIQSQIV SFYLMKFENL
KDDDQRIQRS MDTIKEDMLD KLLNTSSSKR
DDFLKLIQIP VNDLQVQRKA INELFKVMND
LSPRSNLRKR KRSQNLFRGR RASK

```

Specifications

Our [Abpromise guarantee](#) covers the use of **ab93910** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications SDS-PAGE

Purity > 95 % SDS-PAGE.
Purified by Ion-exchange chromatography.

Form Lyophilised

Additional notes Reconstituted protein is stable for at least 3 months when stored in working aliquots with a carrier protein at -20°C. Avoid repeated freeze/thaw cycles.

Preparation and Storage

Stability and Storage Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.

Preservative: None

Constituents: 10% Trehalose, 1X PBS

Reconstitution Reconstitute with sterile phosphate-buffered saline containing at least 0.1% carrier protein.

General Info

Function	Produced by lymphocytes activated by specific antigens or mitogens. IFN-gamma, in addition to having antiviral activity, has important immunoregulatory functions. It is a potent activator of macrophages, it has antiproliferative effects on transformed cells and it can potentiate the antiviral and antitumor effects of the type I interferons.
Tissue specificity	Released primarily from activated T lymphocytes.
Involvement in disease	In Caucasians, genetic variation in IFNG is associated with the risk of aplastic anemia (AA) [MIM:609135]. AA is a rare disease in which the reduction of the circulating blood cells results from damage to the stem cell pool in bone marrow. In most patients, the stem cell lesion is caused by an autoimmune attack. T-lymphocytes, activated by an endogenous or exogenous, and most often unknown antigenic stimulus, secrete cytokines, including IFN-gamma, which would in turn be able to suppress hematopoiesis.
Sequence similarities	Belongs to the type II (or gamma) interferon family.
Post-translational modifications	Proteolytic processing produces C-terminal heterogeneity, with proteins ending alternatively at Gly-150, Met-157 or Gly-161.
Cellular localization	Secreted.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors