

Product datasheet

Recombinant human ERK1 protein ab116536

[1 References](#) [1 Image](#)

Overview

Product name	Recombinant human ERK1 protein
Protein length	Full length protein

Description

Nature	Recombinant
Source	Escherichia coli

Amino Acid Sequence

Accession	P27361
Species	Human
Sequence	MSPILGYWKIKGLVQPTRLLEYLEEKYEEHLYERDEGDKWRNKKFELGL EFPNLPYYIDGDVKLTQSMARIIRYADKHNMLGGCPKERAEISMLEGAVL DIRYGVSRIAYSKDFETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDHVTH PDFMLYDALDVVLYMDPMCLDAFPKLVCFKKRIEAIQIDKYLKSSKYIA WPLQGWQATFGGGDHPPKSDLEVLFGGPLGSAAAAAQGGGGGEPRTGEGV GPGVPGEVEMVKGQPFVGPRTQLQYIGEGAYGMVSSAYDHVRKTRVAI KKISPFHQTYCQRTLREIQILLRFRHENVIGIRDILRASTLEAMRDVYI VQDLMETDLYKLLKSQQLSNDHICYFLYQILRGLKYHSANVLHRDLKPS NLLSNTTCDLKICDFGLARIADPEHDHTGFLTEYVATRWYRAPEIMLNSK GYTKSIDIWSVGCILAEMLSNRPIFPGKHYLDQLNHILGILGSPSQEDLN CIINMKARNYLQSLPSKTKVAWAKLFPKSDSKALDLLDRMLTFNPNKRIT VEEALAHPLYEQYYDPTDEPVAEEPFTFAMELDDLPKERLKELIFQETAR FQPGVLEAP
Molecular weight	70 kDa including tags
Amino acids	2 to 379

Specifications

Our [Abpromise guarantee](#) covers the use of **ab116536** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Biological activity	Specific activity: 379.25 units/ml; 621.73 Units/mg
Applications	SDS-PAGE

	Functional Studies
	Mass Spectrometry
Mass spectrometry	MALDI-TOF-TOF
Purity	> 80 % Proprietary Purification.
Form	Liquid
Additional notes	ab116536 was alkylated, digested with trypsin and the mass of the resultant peptides determined by MADLI-TOF/TOF. The peptides obtained gave 27% sequence coverage of ERK1. Theoretical pI: 6.06


Preparation and Storage

Stability and Storage	Shipped on Dry Ice. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle. pH: 7.50 Constituents: 0.002% Brij, 0.012% Benzamidine, 0.003% EGTA, 0.003% PMSF, 0.1% Beta mercaptoethanol, 0.79% Tris HCl, 9.24% Sucrose, 0.88% Sodium chloride This product is an active protein and may elicit a biological response in vivo, handle with caution.
------------------------------	---

General Info

Function	Involved in both the initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors such as ELK-1. Phosphorylates EIF4EBP1; required for initiation of translation. Phosphorylates microtubule-associated protein 2 (MAP2). Phosphorylates SPZ1 (By similarity). Phosphorylates heat shock factor protein 4 (HSF4).
Sequence similarities	Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily. Contains 1 protein kinase domain.
Domain	The TXY motif contains the threonine and tyrosine residues whose phosphorylation activates the MAP kinases.
Post-translational modifications	Dually phosphorylated on Thr-202 and Tyr-204, which activates the enzyme. Dephosphorylated by PTPRJ at Tyr-204.

Images



12% SDS-PAGE gel showing ab116536.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors