

Product datasheet

Recombinant human NFkB Inducing Kinase NIK protein ab105210

[2 Images](#)

Overview

Product name	Recombinant human NFkB Inducing Kinase NIK protein
Protein length	Protein fragment

Description

Nature	Recombinant
Source	Baculovirus infected Sf9 cells

Amino Acid Sequence

Accession	Q99558
Species	Human
Molecular weight	108 kDa including tags
Amino acids	325 to 947

Specifications

Our [Abpromise guarantee](#) covers the use of **ab105210** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Biological activity The Specific activity of ab105210 was determined to be 8 nmol/min/mg.

Applications Western blot
Functional Studies
SDS-PAGE

Purity > 85 % SDS-PAGE.
Purity was determined to be >85% by densitometry.

Form Liquid

Additional notes [ab64311](#) (Myelin Basic Protein protein) can be utilized as a substrate for assessing kinase activity

Preparation and Storage

Stability and Storage

Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles.

Preservative: 0% None

Constituents: 25% Glycerol, 50mM Tris HCl, 150mM Sodium chloride, 10mM Glutathione, 0.25mM DTT, 0.1mM EDTA, 0.1mM PMSF, pH 7.5

This product is an active protein and may elicit a biological response in vivo, handle with caution.

General Info

Function

Lymphotoxin beta-activated kinase which seems to be exclusively involved in the activation of NF-kappa-B and its transcriptional activity. Promotes proteolytic processing of NFKB2/P100, which leads to activation of NF-kappa-B via the non-canonical pathway. Could act in a receptor-selective manner.

Tissue specificity

Weakly expressed in testis, small intestine, spleen, thymus, peripheral blood leukocytes, prostate, ovary and colon.

Sequence similarities

Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase kinase subfamily.

Contains 1 protein kinase domain.

Post-translational modifications


Autophosphorylated. Phosphorylation at Thr-559 is required to activate its kinase activity and 'Lys-63'-linked polyubiquitination. Phosphorylated by CHUK/IKKA leading to MAP3K14 destabilization.

Ubiquitinated. Undergoes both 'Lys-48'- and 'Lys-63'-linked polyubiquitination. 'Lys-48'-linked polyubiquitination leads to its degradation by the proteasome, while 'Lys-63'-linked polyubiquitination stabilizes and activates it.


Cellular localization

Cytoplasm.

Images



The Specific activity of ab105210 was determined to be 8 nmol/min/mg.



SDS-PAGE showing ab105210 at approximately 108kDa.

SDS-PAGE - NFkB Inducing Kinase NIK protein (ab105210)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors