

Product datasheet

Anti-SEN3 antibody - Aminoterminal end ab71677

1 References 2 Images

Overview

Product name	Anti-SEN3 antibody - Aminoterminal end
Description	Rabbit polyclonal to SEN3 - Aminoterminal end
Host species	Rabbit
Tested applications	Suitable for: ELISA, WB, IHC-P
Species reactivity	Reacts with: Human
Immunogen	KLH-conjugated synthetic peptide, from the N-terminal region of Human SEN3
Positive control	Saos-2 cell lysate; Human breast carcinoma tissue

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Preservative: 0.09% Sodium Azide Constituents: PBS
Purity	Protein G purified
Purification notes	The antibody was purified through a protein G column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab71677** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ELISA		1/1000.
WB		1/100 - 1/500. Predicted molecular weight: 65 kDa.

Application	Abreviews	Notes
IHC-P		1/50 - 1/100.

Target

Function

Protease that releases SUMO2 and SUMO3 monomers from sumoylated substrates, but has only weak activity against SUMO1 conjugates. Deconjugates SUMO2 from MEF2D, which increases its transcriptional activation capability. Deconjugates SUMO2 and SUMO3 from CDCA8. Redox sensor that, when redistributed into nucleoplasm, can act as an effector to enhance HIF1A transcriptional activity by desumoylating EP300. Required for rRNA processing through deconjugation of SUMO2 and SUMO3 from nucleophosmin, NPM1. Plays a role in the regulation of sumoylation status of ZNF148. Functions as a component of the Five Friends of Methylated CHTOP (5FMC) complex; the 5FMC complex is recruited to ZNF148 by methylated CHTOP, leading to desumoylation of ZNF148 and subsequent transactivation of ZNF148 target genes.


Sequence similarities

Belongs to the peptidase C48 family.

Cellular localization

Cytoplasm. Nucleus > nucleolus. Nucleus > nucleoplasm. Mainly found in the nucleoplasm, with low levels detected in the cytoplasmic and chromatin fractions (By similarity). Redistributes between the nucleolus and the nucleoplasm in response to mild oxidative stress.


Images



Anti-SENP3 antibody - Aminoterminal end (ab71677) at 1/100 dilution + Saos-2 cell lysate at 12.5 µg

Predicted band size: 65 kDa

Observed band size: 65 kDa



Analysis of SENP3 expression on formalin-fixed and paraffin-embedded human breast carcinoma tissue using 1:50 ab71677 (peroxidase-conjugated secondary antibody, AEC staining).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - SENP3 antibody - Aminoterminal end (ab71677)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors