


Product datasheet

Anti-STAT5α (phospho Y694) antibody [6.F1] ab106095

★★★★☆ 1 Abreviews 3 Images

Overview

Product name	Anti-STAT5a (phospho Y694) antibody [6.F1]
Description	Mouse monoclonal [6.F1] to STAT5a (phospho Y694)
Host species	Mouse
Specificity	ab106095 is specific for STAT5a protein phosphorylated at Y694.
Tested applications	Suitable for: WB, IHC-P, ICC/IF
Species reactivity	Reacts with: Mouse, Human Predicted to work with: Rat 
Immunogen	A synthetic peptide corresponding to residues surrounding Y694 of Mouse STAT5a protein.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 0.42% Potassium phosphate, 0.88% Sodium chloride
Purity	Protein A purified
Purification notes	NK92 cell lysate; Human breast tissue; 3T3 cells.
Clonality	Monoclonal
Clone number	6.F1
Isotype	IgG1

Applications

Our [Abpromise guarantee](#) covers the use of **ab106095** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
WB		1/500 - 1/2000. Predicted molecular weight: 91 kDa.

Application	Abreviews	Notes
IHC-P	★★★★☆	Use a concentration of 20 - 40 µg/ml.
ICC/IF		1/50 - 1/1000.

Target

Function	Carries out a dual function: signal transduction and activation of transcription. Binds to the GAS element and activates PRL-induced transcription.
Sequence similarities	Belongs to the transcription factor STAT family. Contains 1 SH2 domain.
Post-translational modifications	Tyrosine phosphorylated in response to IL-2, IL-3, IL-7, IL-15, GM-CSF, growth hormone, prolactin, erythropoietin and thrombopoietin. Tyrosine phosphorylation is required for DNA-binding activity and dimerization. Serine phosphorylation is also required for maximal transcriptional activity.
Cellular localization	Cytoplasm. Nucleus. Translocated into the nucleus in response to phosphorylation.

Images



All lanes : Anti-STAT5a (phospho Y694) antibody [6.F1] (ab106095) at 1/1000 dilution (overnight at 4°C)


Lane 1 : NK92 cell lysate, untreated

Lane 2 : NK92 cell lysate, treated with 1Ku of IL2 (30min)

Secondary


All lanes : Gt-a-Mouse IgG DyLight™800 conjugated at 1/20000 dilution

Predicted band size: 91 kDa



ab106095 at 20µg/ml staining STAT5a in Formalin-fixed, Paraffin-embedded Human breast tissue by Immunohistochemistry. The image shows the localization of antibody as the precipitated red signal, with a hematoxylin purple nuclear counterstain.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STAT5a (phospho Y694) antibody [6.F1] (ab106095)



ab106095 staining STAT5a in 3T3 cells by Immunofluorescence (red). Green staining represents tubulin.

Immunocytochemistry/ Immunofluorescence - Anti-STAT5a (phospho Y694) antibody [6.F1] (ab106095)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors