


Product datasheet

Anti-TIRAP antibody ab17218

★★★★★ 2 Abreviews 4 References 4 Images

Overview

| | |
|----------------------------|---|
| Product name | Anti-TIRAP antibody |
| Description | Rabbit polyclonal to TIRAP |
| Tested applications | Suitable for: WB, ICC/IF, IHC-P |
| Species reactivity | Reacts with: Mouse, Rat, Human Predicted to work with: Rhesus monkey  |
| Immunogen | Synthetic peptide. 15 amino acids near the C-terminus of mouse TIRAP (GenBank accession no.AAL05628). |
| Positive control | Human heart cell lysates |

Properties

| | |
|-----------------------------|---|
| Form | Liquid |
| Storage instructions | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C. Avoid freeze / thaw cycle. Stable for 12 months at -20°C. |
| Storage buffer | Preservative: 0.02% Sodium Azide Constituents: PBS |
| Purity | Immunogen affinity purified |
| Purification notes | ab17218 is purified by ion exchange chromatography. TIRAP Antibody is affinity chromatography purified via peptide column. |
| Clonality | Polyclonal |
| Isotype | IgG |

Applications

Our [Abpromise guarantee](#) covers the use of **ab17218** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|-------------|-----------|--|
| WB | | Use a concentration of 0.5 - 2 µg/ml. Detects a band of approximately 34 kDa.Can be blocked with Mouse TIRAP peptide (ab39884) . |

| Application | Abreviews | Notes |
|-------------|-----------|--|
| ICC/IF | ★★★★☆ | Use at an assay dependent concentration. |
| IHC-P | ★★★★★ | Use a concentration of 2 µg/ml. |

Target


Relevance

The Toll-like receptor (TLR) family in mammals comprises a family of transmembrane proteins characterized by multiple copies of leucine rich repeats in the extracellular domain and IL-1 receptor motif in the cytoplasmic domain. Like its counterparts in *Drosophila*, TLRs signal through adaptor molecules. The TLR family is a phylogenetically conserved mediator of innate immunity that is essential for microbial recognition. Ten human homologs of TLRs (TLR1-10) have been described. TIRAP (TIR domain-containing adaptor protein) is an adaptor protein used by TLR4. Blocking TIRAP inhibits TLR4-mediated signaling events, including DC maturation and cytokine production. Function: Adapter involved in TLR2 and TLR4 signaling pathways in the innate immune response. Acts via IRAK2 and TRAF-6, leading to the activation of NF-kappa-B, MAPK1, MAPK3 and JNK, and resulting in cytokine secretion and the inflammatory response. Positively regulates the production of TNF-alpha and interleukin-6. Tissue specificity: Highly expressed in liver, kidney, spleen, skeletal muscle and heart. Also detected in peripheral blood leukocytes, lung, placenta, small intestine, thymus, colon and brain. Post-translational modification: Phosphorylated by IRAK1 and IRAK4. Also phosphorylated by BTK. Sequence similarities: Contains 1 TIR domain.


Cellular localization

Cytoplasmic

Images



Western Blot of human heart lysate labeling TIRAP with Anti-TIRAP antibody (ab17218) at (1) 1 and (2) 2µg/ml.



Western blot - TIRAP antibody (ab17218)

Lane 1 : Anti-TIRAP antibody (ab17218) at 0.5 µg/ml

Lane 2 : Anti-TIRAP antibody (ab17218) at 1 µg/ml

Lane 3 : Anti-TIRAP antibody (ab17218) at 2 µg/ml

Lane 1 : heart cell lysate

Lane 2 : heart cell lysate


Lane 3 : heart cell lysate

Observed band size : 35 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - TIRAP antibody (ab17218)

ab17218 at 2µg/ml staining TIRAP in human heart cells by IHC



Immunocytochemistry/ Immunofluorescence - TIRAP antibody (ab17218)

This image is courtesy of an Anonymous Abreview.

ab17218 staining TIRAP in a PMA-treated Human THP-1 cell by Immunocytochemistry/ Immunofluorescence. Cells were fixed in paraformaldehyde for 30 minutes and permeabilized in 0.1% Triton X-100 prior to blocking in 5% serum for 1 hour at 20°C. The primary antibody was diluted 1/200 and incubated with the sample for 1 hour at 20°C. The secondary antibody was Alexa Fluor® 488-conjugated goat anti-rabbit polyclonal, diluted 1/500 (green). Other stains used were Alexa Fluor 595-conjugated wheat germ agglutinin (red) and Nuclei were stained with Hoechst 33258 (blue)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors