

Product datasheet

Anti-TRIP15 antibody ab4537

[1 References](#) [2 Images](#)

Overview

Product name	Anti-TRIP15 antibody
Description	Goat polyclonal to TRIP15
Tested applications	Suitable for: IHC-P, WB
Species reactivity	Reacts with: Mouse, Human Predicted to work with: Rat
Immunogen	Synthetic peptide: LNSLNQAVVSKLA, corresponding to C terminal amino acids 421-443 of Human TRIP15. Run BLAST with ExPASy Run BLAST with NCBI
Positive control	Purchase matching WB positive control: Recombinant TRIP15 protein >
General notes	GenBank Accession Number – NP_004227. Gene Ontology terms - protein binding; signal transducer activity; signal transduction; transcription from Pol II promoter; cytoplasm; signalosome complex. LocusLink ID - 9318 (human); 12848 (mouse); 261736 (rat).

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
Storage buffer	Preservative: 0.02% Sodium Azide Constituents: 0.5% BSA, Tris buffered saline
Purity	Immunogen affinity purified
Purification notes	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab4537** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		Use a concentration of 3 - 5 µg/ml.
WB		Use a concentration of 0.03 - 3 µg/ml. Predicted molecular weight: 51 kDa. Can be blocked with Human TRIP15 peptide (ab23046) .

Target


Function	Essential component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, IκappaBα/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively. Involved in early stage of neuronal differentiation via its interaction with NIF3L1.
Sequence similarities	Belongs to the CSN2 family. Contains 1 PCI domain.
Post-translational modifications	Phosphorylated by CK2 and PKD kinases.
Cellular localization	Cytoplasm. Nucleus.

Images



Anti-TRIP15 antibody (ab4537) at 0.03 µg/ml
+ NIH-3T3 cell lysate at 30 µg
Developed using the ECL technique

Predicted band size : 51 kDa



ab4537 (3-5 µg/ml) staining TRIP15 in paraffin-embedded sections of Human thyroid Gland.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - TRIP15 antibody (ab4537)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors