

## Product datasheet

### XIAP peptide ab38275

#### Overview

---

**Product name** XIAP peptide

#### Description

---

**Nature** Synthetic

#### Specifications

---

Our [Abpromise guarantee](#) covers the use of **ab38275** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

**Applications** Blocking - Blocking peptide for Anti-XIAP antibody ([ab2541](#))  
Western blot

**Form** Liquid

#### Preparation and Storage

---

**Stability and Storage** Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.  
Preservative: 0.02% Thimerosal (merthiolate)  
Constituents: 0.1% BSA, PBS, pH 7.2

#### General Info

---

**Function** Apoptotic suppressor. Has E3 ubiquitin-protein ligase activity. Mediates the proteasomal degradation of target proteins, such as caspase-3, SMAC or AIFM1. Inhibitor of caspase-3, -7 and -9. Mediates activation of MAP3K7/TAK1, leading to the activation of NF-kappa-B.

**Tissue specificity** Ubiquitous, except peripheral blood leukocytes.

**Involvement in disease** Defects in XIAP are the cause of lymphoproliferative syndrome X-linked type 2 (XLP2) [MIM:300635]. XLP is a rare immunodeficiency characterized by extreme susceptibility to infection with Epstein-Barr virus (EBV). Symptoms include severe or fatal mononucleosis, acquired hypogammaglobulinemia, pancytopenia and malignant lymphoma.

**Sequence similarities** Belongs to the IAP family.  
Contains 3 BIR repeats.  
Contains 1 RING-type zinc finger.

<b>Domain</b>	The first BIR domain is involved in interaction with TAB1/MAP3K7IP1 and is important for dimerization. The second BIR domain is sufficient to inhibit caspase-3 and caspase-7, while the third BIR is involved in caspase-9 inhibition. The interactions with SMAC and PRSS25 are mediated by the second and third BIR domains.
<b>Post-translational modifications</b>	Ubiquitinated and degraded by the proteasome in apoptotic cells. Phosphorylation by PKB/AKT protects XIAP against ubiquitination and protects the protein against proteasomal degradation.
<b>Cellular localization</b>	Cytoplasm.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors