

Product datasheet

Anti-ZP3 antibody ab48895

★★★★☆ 3 Abreviews 1 Image

Overview

Product name	Anti-ZP3 antibody
Description	Rabbit polyclonal to ZP3
Host species	Rabbit
Tested applications	Suitable for: ICC, WB
Species reactivity	Reacts with: Human Predicted to work with: Cow
Immunogen	A region within synthetic peptide: QWSRSASRNR RHVTEEADVTVGGLIFLDRR GDHEVEQWAL PSDTSVLLG, corresponding to internal sequence amino acids 342-391 of Human ZP3 Run BLAST with ExPASy Run BLAST with NCBI
Positive control	HepG2 cell lysate

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Preservative: None Constituents: 2% Sucrose, PBS
Purity	Protein A purified
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab48895** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
ICC	★★★★☆	Use at an assay dependent concentration.

Application	Abreviews	Notes
WB	★★★★★	Use a concentration of 0.6 µg/ml. Detects a band of approximately 47 kDa (predicted molecular weight: 47 kDa). Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.

Target

Function	The mammalian zona pellucida, which mediates species-specific sperm binding, induction of the acrosome reaction and prevents post-fertilization polyspermy, is composed of three to four glycoproteins, ZP1, ZP2, ZP3, and ZP4. ZP3 is essential for sperm binding and zona matrix formation.
Tissue specificity	Oocytes.
Sequence similarities	Belongs to the ZP domain family. ZPC subfamily. Contains 1 ZP domain.
Domain	The ZP domain is involved in the polymerization of the ZP proteins to form the zona pellucida.
Post-translational modifications	Proteolytically cleaved before the transmembrane segment to yield the secreted ectodomain incorporated in the zona pellucida. N-glycosylated. O-glycosylated; removal of O-linked glycans may play an important role in the post-fertilization block to polyspermy.
Cellular localization	Secreted > extracellular space > extracellular matrix and Cell membrane.

Images



Western blot - Anti-ZP3 antibody (ab48895)

Anti-ZP3 antibody (ab48895) at 0.6 µg/ml +
HepG2 cell lysate at 10 µg

Secondary

HRP conjugated anti-Rabbit IgG at 1/50000
dilution

Developed using the ECL technique.

Predicted band size: 47 kDa

Observed band size: 45 kDa

Diluted in 5% skim milk / PBS buffer, gel
concentration 12%

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors