

Product datasheet

Anti-MHC Class 1 H2 Db antibody [28-14-8] (Biotin) ab25237

1 References

Overview

Product name	Anti-MHC Class 1 H2 Db antibody [28-14-8] (Biotin)
Description	Mouse monoclonal [28-14-8] to MHC Class 1 H2 Db (Biotin)
Conjugation	Biotin
Specificity	This antibody binds to the a3 domain of H-2Db in the presence or absence of β 2 microglobulin. It cross reacts with the a3 domain of H-2Ld, but not Kd or Dd, and with H-2Dq and/or Lq.
Tested applications	Suitable for: Flow Cyt, IHC-Fr
Species reactivity	Reacts with: Mouse
Immunogen	C3H.SW mouse splenocytes.
Epitope	This antibody recognises an epitope in the a3 domain of H-2Db.
General notes	This antibody has been shown to block H-2Ld-specific and H-2Ld-restricted antigen-specific lysis of target cells by cytotoxic T lymphocytes, but it does not block recognition of H-2Ld-positive target cells by Ly-6G2-positive NK cells. This antibody has been shown to be useful in studies of complement mediated cytotoxicity.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.2% Sodium azide Constituent: PBS
Purity	IgG fraction
Primary antibody notes	This antibody has been shown to block H-2Ld-specific and H-2Ld-restricted antigen-specific lysis of target cells by cytotoxic T lymphocytes, but it does not block recognition of H-2Ld-positive target cells by Ly-6G2-positive NK cells. This antibody has been shown to be useful in studies of complement mediated cytotoxicity.
Clonality	Monoclonal
Clone number	28-14-8

Isotype	IgG2a
Light chain type	kappa

Applications

Our [Abpromise guarantee](#) covers the use of **ab25237** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		Use 1µg for 10 ⁶ cells. ab18454 -Mouse monoclonal IgG2a, is suitable for use as an isotype control with this antibody.
IHC-Fr		Use at an assay dependent concentration.

Target

Relevance

In the mouse the MHC Class 1 loci are called H2K, H2L and H2D. The equivalent loci in the human MHC are HLA (Human Leukocyte Antigen). MHC class I molecules are heterodimers, consisting of a single transmembrane polypeptide chain (the α-chain) which is highly polymorphic, and the invariant β2 microglobulin (which is encoded elsewhere, not in the MHC). MHC class I molecules are found on almost every nucleated cell of the body. Their major function is to present peptide fragments derived from antigens to cytotoxic T cells.

Cellular localization

Cell Membrane; single pass type I membrane protein

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors