

Product datasheet

Anti-ADAM20 antibody - Catalytic domain ab28295

1 Image

Overview

Product name	Anti-ADAM20 antibody - Catalytic domain
Description	Rabbit polyclonal to ADAM20 - Catalytic domain
Tested applications	Suitable for: WB
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide based on the catalytic domain of human ADAM20. Read Abcam's proprietary immunogen policy (Peptide available as ab41221 .)

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
Storage buffer	Preservative: 0.05% Sodium Azide Constituents: 50% Glycerol
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab28295** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		

Application notes


WB: A recommended starting concentration for Western Blots is 1:1000 when using colorimetric substrates such as BCIP/NBT, and 1:5000 for chemiluminescent substrates. Predicted molecular weight: 82 kDa. Higher concentrations of antibody may be needed for samples from more distantly related species. EDTA/EGTA treatment of tissues or lysates is required to see latent zymogen. This product recognizes bands of 95 kDa, 84 kDa (major band), and breakdown

products at 50 kDa, 34 kDa in reduced Western Blots of cell lysates. Dilution optimised using Chromogenic detection. Not yet tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Target

Function	May be involved in sperm maturation and/or fertilization.
Tissue specificity	Testis specific.
Sequence similarities	Contains 1 disintegrin domain. Contains 1 EGF-like domain. Contains 1 peptidase M12B domain.
Domain	A tripeptide motif (VGE) within disintegrin-like domain could be involved in the binding to egg integrin receptor and thus could mediate sperm/egg binding. The cysteine-rich domain encodes putative cell-fusion peptides, which could be involved in sperm-egg fusion. The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.
Post-translational modifications	Has no obvious cleavage site for furin endopeptidase, suggesting that the proteolytic processing is regulated.
Cellular localization	Membrane.

Images



Western blot - ADAM20 antibody - Catalytic domain (ab28295)

All lanes : Anti-ADAM20 antibody - Catalytic domain (ab28295) at 1/1000 dilution

Lane 1 : Mouse Testis Lysate

Lane 2 : Mouse Epididymis Lysate

Lysates/proteins at 20 µl per lane.

Predicted band size : 82 kDa

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors