

Product datasheet

Anti-CD4 antibody [YTS 177.9] (Phycoerythrin) ab33437

5 References

Overview

Product name	Anti-CD4 antibody [YTS 177.9] (Phycoerythrin)
Description	Rat monoclonal [YTS 177.9] to CD4 (Phycoerythrin)
Host species	Rat
Conjugation	Phycoerythrin. Ex: 488nm, Em: 575nm
Specificity	This antibody reacts with the CD4 antigen, non polymorphic epitope. The antibody reacts with CD4 transfectants and blocks MHCII dependant T cell responses in vitro and in vivo and induces tolerance.
Tested applications	Suitable for: Flow Cyt
Species reactivity	Reacts with: Mouse
Immunogen	Tissue/ cell preparation: Mouse spleen cells.

Properties

Form	Lyophilised:Reconstitute with 1ml distilled water.
Storage instructions	Store at +4°C in the dark. Do not freeze.
Storage buffer	Preservative: 0.09% Sodium Azide Constituents: 1.0% BSA, PBS, pH 7.2
Purity	Protein G purified
Clonality	Monoclonal
Clone number	YTS 177.9
Myeloma	Y3/Ag1.2.3
Isotype	IgG2a

Applications

Our [Abpromise guarantee](#) covers the use of **ab33437** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
-------------	-----------	-------

Flow Cyt

Application notes

Flow Cyt: Use 10µl for 10⁶ cells or 100µl of whole blood.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

Target

Function

Accessory protein for MHC class-II antigen/T-cell receptor interaction. May regulate T-cell activation. Induces the aggregation of lipid rafts.

Sequence similarities

Contains 3 Ig-like C2-type (immunoglobulin-like) domains.

Contains 1 Ig-like V-type (immunoglobulin-like) domain.

Post-translational modifications

Palmitoylation and association with LCK contribute to the enrichment of CD4 in lipid rafts.

Cellular localization

Cell membrane. Localizes to lipid rafts. Removed from plasma membrane by HIV-1 Nef protein that increases clathrin-dependent endocytosis of this antigen to target it to lysosomal degradation. Cell surface expression is also down-modulated by HIV-1 Envelope glycoprotein gp160 that interacts with, and sequesters CD4 in the endoplasmic reticulum.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors