

Product datasheet

Anti-S6K1 antibody ab36864

2 References 3 Images

Overview

Product name	Anti-S6K1 antibody
Description	Rabbit polyclonal to S6K1
Host species	Rabbit
Tested applications	Suitable for: ICC/IF, WB
Species reactivity	Reacts with: Human
Immunogen	A 16 amino acid peptide from near the C terminus of human RPS6KB1.
Positive control	Jurkat cell lysate.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C.
Storage buffer	Preservative: 0.02% Sodium Azide Constituents: PBS
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab36864** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		Use a concentration of 10 µg/ml.
WB		Use a concentration of 5 - 10 µg/ml. Detects a band of approximately 74 kDa (predicted molecular weight: 59 kDa).

Target

Function

Acts to integrate nutrient and growth factor signals in regulation of protein synthesis, cell proliferation, cell growth, cell cycle progression and cell survival. Downstream effector of the mTOR signaling pathway. Phosphorylates specifically ribosomal protein S6 in response to insulin or several classes of mitogens. During translation initiation, the inactive form associates with the eIF-3 complex under conditions of nutrient depletion. Mitogenic stimulation leads to phosphorylation and dissociation from the eIF-3 complex and the free activated form can phosphorylate other translational targets including EIF4B. Promotes protein synthesis by phosphorylating PDCD4 at 'Ser-67' and targeting it for degradation. Phosphorylates RICTOR leading to regulation of mammalian target of rapamycin complex 2 (mTORC2) signaling; probably phosphorylates RICTOR at 'Thr-1135'. Phosphorylates IRS1 at multiple serine residues coupled with insulin resistance; probably phosphorylates IRS1 at 'Ser-270'. Required for TNF-alpha induced IRS-1 degradation. Phosphorylates EEF2K in response to IGF1 and inhibits EEF2K activity. Phosphorylates BAD at 'Ser-99' in response to IGF1 leading to BAD inactivation and inhibition of BAD-induced apoptosis. Phosphorylates mitochondrial RMP leading to dissociation of a RMP:PPP1CC complex; probably phosphorylates RMP at 'Ser-99'. The free mitochondrial PPP1CC can dephosphorylate RPS6KB1 at Thr-412 which is proposed to be a negative feed back mechanism for the RPS6KB1 antiapoptotic function. Phosphorylates GSK3B at 'Ser-9' under conditions leading to loss of the TSC1-TSC2 complex. Phosphorylates POLDIP3.

Tissue specificity

Widely expressed.

Sequence similarities

Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. S6 kinase subfamily.

Contains 1 AGC-kinase C-terminal domain.

Contains 1 protein kinase domain.

Domain

The autoinhibitory domain is believed to block phosphorylation within the AGC-kinase C-terminal domain and the activation loop.

The TOS (TOR signaling) motif is essential for activation by mTORC1.


Post-translational modifications

Phosphorylation at Thr-412 is regulated by mTORC1. The phosphorylation at this site is maintained by an agonist-dependent autophosphorylation mechanism.

Cellular localization


Cytoplasm; Nucleus. Cytoplasm and Cell junction > synapse > synaptosome. Mitochondrion outer membrane.

Images



Immunocytochemistry/ Immunofluorescence - Anti-S6K1 antibody (ab36864)

Immunofluorescence of S6K1 in Jurkat cells with ab36864 at 20 ug/mL.



Western blot - S6K antibody (ab36864)

Lane 1 : Anti-S6K1 antibody (ab36864) at 2.5 μ g/ml

Lane 2 : Anti-S6K1 antibody (ab36864) at 5 μ g/ml

Lane 3 : Anti-S6K1 antibody (ab36864) at 10 μ g/ml

All lanes : Jurkat cell lysate

Lysates/proteins at 15 μ g/ml per lane.

Secondary

All lanes : anti-rabbit IgG HRP at 1/10000 dilution

Predicted band size: 59 kDa

Observed band size: 74 kDa

Lanes read from left to right, 1, 2 and 3.



ab36864 at 10µg/ml staining S6K in Jurkat cells by Immunocytochemistry.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors