# abcam

## Product datasheet

## ILl alpha Human ELISA Kit ab46051

## 1 Image

Overview

**Product name** 

IL1 alpha Human ELISA Kit

**Detection method** 

Colorimetric

Precision

				ii iti a assay
Sample	n	Mean	SD	CV%
Α	22	459.6pg/ml	13	2.8%
В	21	72.6pg/ml	3.6	4.3%

Inter-assay

Intra-assav

Sample	n	Mean	SD	CV%
Α	8	259.9pg/ml	15	5.8%
В	8	58.6pg/ml	4.3	7.3%

Sample type Cell culture supernatant, Serum, Plasma

Assay type Sandwich (quantitative)

Sensitivity < 10 pg/ml

**Range** 31.2 pg/ml - 1000 pg/ml

Recovery 95.3 %

Sample specific recovery

Sample type	Average %	Range
Serum	95.3	31.25pg/ml - 1000pg/ml

Assay time 3h 45m

**Assay duration** Multiple steps standard assay

Species reactivity Reacts with: Human

Product overview The Human IL1 alpha ELISA is to be used for the in-vitro quantitative determination of

interleukin-1 alpha (IL1 alpha) in human serum, plasma, buffered solutions or cell culture

1

medium. The assay will recognize both natural and recombinant human IL1 alpha. This kit has been configured for research use only and is not to be used in diagnostic procedures.

#### Principal:

The IL1 alpha Kit is a solid phase sandwich Enzyme Linked-Immuno-Sorbent Assay (ELISA). A monoclonal antibody specific for IL1 alpha has been coated onto the wells of the microtiter strips provided. Samples, including standards of known IL1 alpha concentrations and unknowns are pipetted into these wells. During the first incubation, the IL1 alpha antigen is added to wells. After washing, a biotinylated monoclonal antibody specific for IL1 alpha is incubated. Then the enzyme (streptavidin-peroxydase) is added. After incubation and washing to remove all unbound enzyme, a substrate solution which acts on the bound enzyme is added to induce a coloured reaction product. The intensity of this coloured product is directly proportional to the concentration of IL1 alpha present in the samples.

Recognizes both natural and recombinant human IL1 alpha

Range: 31.2 pg/ml - 1000 pg/ml

Sensitivity <10 pg/ml

Notes Safety Notes:

For research use only.

The human blood components included in this kit have been tested and found non reactive for HBsAg and anti-HIV. Nevertheless, no known method can offer complete assurance that human blood derivatives will not transmit hepatitis, AIDS or other infections. Therefore, handling of reagents, serum or plasma specimens should be in accordance with local safety procedures, e.g. CDC/NIH Health manual: "Biosafety in Microbiological and Biomedical Laboratories" 1984.

Avoid any skin contact with  $H_2SO_4$  and TMB. In case of contact, wash thoroughly with water.

Do not eat, drink, smoke or apply cosmetics where kit reagents are used.

Do not pipette by mouth.

The stop solution in this kit contains 9.8% sulphuric acid.

**Tested applications** 

Suitable for: ELISA

**Platform** 

Microplate

**Properties** 

Storage instructions

Store at +4°C. Please refer to protocols.

3	
Components	
10X Standard Diluent Buffer	1 x 25ml
Biotinylated Antibody Diluent	1 x 13ml
HRP Diluent	1 x 23ml

Components		
Stop Reagent		2 x 11ml
Chromogen TMB Substrate Solution		1 x 24ml
200X Wash Buffer		2 x 10ml
Biotinylated anti-IL1a		2 x 0.4ml
Streptavidin-HRP		4 x 5µl
IL-1a Microplate (12 x 8 well strips)		8 x unit
IL-1a Standard (Lyophilized)		4 x 1vial
IL1a Control		4 x 1vial
Relevance	IL1 alpha is a member of the interleukin 1 cytokine family. It is a pleiotropic cytokine involved in various immune responses, inflammatory processes, and hematopoiesis. IL1 alpha is produced by monocytes and macrophages as a proprotein, which is proteolytically processed and released in response to cell injury, and thus induces apoptosis. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. It has been suggested that the polymorphism of these genes is associated with rheumatoid arthritis and Alzheimer's disease.	
Cellular localization	Secreted	


## **Applications**

Our Abpromise guarantee covers the use of **ab46051** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ELISA		Use at an assay dependent dilution.

## **Images**



Typical IL1 alpha standard curve ranging from 1000 to 31.25 pg/mL.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="http://www.abcam.com/abpromise">http://www.abcam.com/abpromise</a> or contact our technical team.

#### Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors