

## Product datasheet

# Anti-SF2 antibody ab50565

1 Image

### Overview

<b>Product name</b>	Anti-SF2 antibody
<b>Description</b>	Rabbit polyclonal to SF2
<b>Specificity</b>	This antibody reacts with SF2
<b>Tested applications</b>	<b>Suitable for:</b> WB, ELISA
<b>Species reactivity</b>	<b>Reacts with:</b> Human <b>Predicted to work with:</b> Mouse, Rat, Cow, Dog, Pig, Zebrafish
<b>Immunogen</b>	A region within synthetic peptide: EFVRKEDMTY AVRKLDNTKF RSHEGETAYI RVKVDGPRSP SYGRSRSRSR, corresponding to amino acids 161-210 of Human SF2 <a href="#">Run BLAST with ExPASy</a> <a href="#">Run BLAST with NCBI</a>
<b>Positive control</b>	HepG2 cell lysate

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
<b>Storage buffer</b>	Preservative: None Constituents: 2% Sucrose, PBS
<b>Purity</b>	Immunogen affinity purified
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG


### Applications

Our [Abpromise guarantee](#) covers the use of **ab50565** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use a concentration of 0.25 µg/ml. Detects a band of approximately 32 kDa (predicted molecular weight: 27 kDa). Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.

Application	Abreviews	Notes
ELISA		1/62500. Titre using peptide based assay: 1/62500.
<hr/>		
<b>Target</b>		
<hr/>		
<b>Function</b>		Plays a role in preventing exon skipping, ensuring the accuracy of splicing and regulating alternative splicing. Interacts with other spliceosomal components, via the RS domains, to form a bridge between the 5'- and 3'-splice site binding components, U1 snRNP and U2AF. Can stimulate binding of U1 snRNP to a 5'-splice site-containing pre-mRNA. Binds to purine-rich RNA sequences, either the octamer, 5'-RGAAGAAC-3' (r=A or G) or the decamers, AGGACAGAGC/AGGACGAAGC. Binds preferentially to the 5'-CGAGGCG-3' motif in vitro. Three copies of the octamer constitute a powerful splicing enhancer in vitro, the ASF/SF2 splicing enhancer (ASE) which can specifically activate ASE-dependent splicing. Isoform ASF-2 and isoform ASF-3 act as splicing repressors.
<b>Sequence similarities</b>		Belongs to the splicing factor SR family. Contains 2 RRM (RNA recognition motif) domains.
<b>Domain</b>		The RRM 2 domain plays an important role in governing both the binding mode and the phosphorylation mechanism of the RS domain by SRPK1. RS domain and RRM 2 are uniquely positioned to initiate a highly directional (C-terminus to N-terminus) phosphorylation reaction in which the RS domain slides through an extended electronegative channel separating the docking groove of SRPK1 and the active site. RRM 2 binds toward the periphery of the active site and guides the directional phosphorylation mechanism. Both the RS domain and an RRM domain are required for nucleocytoplasmic shuttling.
<b>Post-translational modifications</b>		Phosphorylated by CLK1, CLK2, CLK3 and CLK4. Phosphorylated by SRPK1 at multiple serines in its RS domain via a directional (C-terminal to N-terminal) and a dual-track mechanism incorporating both processive phosphorylation (in which the kinase stays attached to the substrate after each round of phosphorylation) and distributive phosphorylation steps (in which the kinase and substrate dissociate after each phosphorylation event). The RS domain of SRSF1 binds to a docking groove in the large lobe of the kinase domain of SRPK1 and this induces certain structural changes in SRPK1 and/or RRM 2 domain of SRSF1, allowing RRM 2 to bind the kinase and initiate phosphorylation. The cycles continue for several phosphorylation steps in a processive manner (steps 1-8) until the last few phosphorylation steps (approximately steps 9-12). During that time, a mechanical stress induces the unfolding of the beta-4 motif in RRM 2, which then docks at the docking groove of SRPK1. This also signals RRM 2 to begin to dissociate, which facilitates SRSF1 dissociation after phosphorylation is completed. Arg-97 is dimethylated, probably to asymmetric dimethylarginine.
<b>Cellular localization</b>		Cytoplasm. Nucleus speckle. In nuclear speckles. Shuttles between the nucleus and the cytoplasm.
<hr/>		
<b>Images</b>		
<hr/>		



Western blot - SF2 antibody (ab50565)

**Lane 1 :**

**Lane 2 :** Anti-SF2 antibody (ab50565) at 0.25  $\mu\text{g/ml}$

**Lane 1 :** MARKER

**Lane 2 :** HepG2 cell lysate at 10  $\mu\text{g}$

**Secondary**

**Lane 2 :** HRP conjugated anti-Rabbit IgG at 1/50000 dilution

**Predicted band size :** 27 kDa

**Observed band size :** 32 kDa

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors