


## Product datasheet

# Anti-beta Catenin (phospho S675) antibody ab58615

1 Image

### Overview

---

<b>Product name</b>	Anti-beta Catenin (phospho S675) antibody
<b>Description</b>	Rabbit polyclonal to beta Catenin (phospho S675)
<b>Specificity</b>	ab58615 reacts specifically with phosphorylated $\beta$ -Catenin at the phosphorylation site of Ser675, not with other non-related phosphospecific peptides or non-phosphorylated $\beta$ -Catenin peptides.
<b>Tested applications</b>	<b>Suitable for:</b> ELISA, IHC-Fr, IP, WB
<b>Species reactivity</b>	<b>Reacts with:</b> Human <b>Predicted to work with:</b> Mouse, Rat, Chicken 
<b>Immunogen</b>	A synthetic peptide of C-terminal portion of human $\beta$ -Catenin at the phosphorylation site of Serine 675.
<b>Positive control</b>	Mouse embryonic cell lysate HEK293 whole cell lysate CCL-226 whole cell lysate
<b>General notes</b>	Store at 2-8 °C for up to 1 year.

### Properties

---

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C.
<b>Storage buffer</b>	Preservative: 0.05% Sodium Azide Constituents: PBS, pH 7.4
<b>Purity</b>	Immunogen affinity purified
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG

### Applications

---

Our [Abpromise guarantee](#) covers the use of **ab58615** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ELISA		Use a concentration of 0.1 - 1 µg/ml.
IHC-Fr		Use a concentration of 1 - 2 µg/ml.
IP		Use 3µg for 10 <sup>5</sup> cells.
WB		Use a concentration of 0.5 - 2 µg/ml. Predicted molecular weight: 88 kDa.

## Target

### Function

Key downstream component of the canonical Wnt signaling pathway. In the absence of Wnt, forms a complex with AXIN1, AXIN2, APC, CSNK1A1 and GSK3B that promotes phosphorylation on N-terminal Ser and Thr residues and ubiquitination of CTNNB1 via BTRC and its subsequent degradation by the proteasome. In the presence of Wnt ligand, CTNNB1 is not ubiquitinated and accumulates in the nucleus, where it acts as a coactivator for transcription factors of the TCF/LEF family, leading to activate Wnt responsive genes.

Involved in the regulation of cell adhesion. The majority of beta-catenin is localized to the cell membrane and is part of E-cadherin/catenin adhesion complexes which are proposed to couple cadherins to the actin cytoskeleton.

### Tissue specificity

Expressed in several hair follicle cell types: basal and peripheral matrix cells, and cells of the outer and inner root sheaths. Expressed in colon.

### Involvement in disease

Defects in CTNNB1 are associated with colorectal cancer (CRC) [MIM:114500].  
 Note=Activating mutations in CTNNB1 have oncogenic activity resulting in tumor development. Somatic mutations are found in various tumor types, including colon cancers, ovarian and prostate carcinomas, hepatoblastoma (HB), hepatocellular carcinoma (HCC). HBs are malignant embryonal tumors mainly affecting young children in the first three years of life.

Defects in CTNNB1 are a cause of pilomatixoma (PTR) [MIM:132600]; a common benign skin tumor.

Defects in CTNNB1 are a cause of medulloblastoma (MDB) [MIM:155255]. MDB is a malignant, invasive embryonal tumor of the cerebellum with a preferential manifestation in children.

Defects in CTNNB1 are a cause of susceptibility to ovarian cancer (OC) [MIM:167000]. Ovarian cancer common malignancy originating from ovarian tissue. Although many histologic types of ovarian neoplasms have been described, epithelial ovarian carcinoma is the most common form. Ovarian cancers are often asymptomatic and the recognized signs and symptoms, even of late-stage disease, are vague. Consequently, most patients are diagnosed with advanced disease.

Note=A chromosomal aberration involving CTNNB1 is found in salivary gland pleiomorphic adenomas, the most common benign epithelial tumors of the salivary gland. Translocation t(3;8) (p21;q12) with PLAG1.

### Sequence similarities

Belongs to the beta-catenin family.  
 Contains 12 ARM repeats.

### Post-translational modifications

Phosphorylation by GSK3B requires prior phosphorylation of Ser-45 by another kinase. Phosphorylation proceeds then from Thr-41 to Ser-37 and Ser-33.

EGF stimulates tyrosine phosphorylation. Phosphorylation on Tyr-654 decreases CDH1 binding and enhances TBP binding.


Ubiquitinated by the SCF(BTRC) E3 ligase complex when phosphorylated by GSK3B, leading to its degradation. Ubiquitinated by a E3 ubiquitin ligase complex containing UBE2D1, SIAH1,

CACYBP/SIP, SKP1, APC and TBL1X, leading to its subsequent proteasomal degradation.

## Cellular localization

Cytoplasm. Nucleus. Cytoplasm > cytoskeleton. Cell junction > adherens junction. Cell junction. Cell membrane. Cytoplasmic when it is unstabilized (high level of phosphorylation) or bound to CDH1. Translocates to the nucleus when it is stabilized (low level of phosphorylation). Interaction with GLIS2 and MUC1 promotes nuclear translocation. Interaction with EMD inhibits nuclear localization.

## Images



Western blot - beta Catenin (phospho S675)  
antibody (ab58615)

**All lanes :** Anti-beta Catenin (phospho S675)  
antibody (ab58615) at 1/200 dilution

**Lane 1 :** (Left): Stimulated CCL-226 whole  
cell lysate, minus immunising peptide

**Lane 2 :** (Right): Stimulated CCL-226 whole  
cell lysate, plus immunising peptide

Lysates/proteins at 15 µg per lane.

**Predicted band size :** 88 kDa

**Observed band size :** 88 kDa

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors