


Product datasheet

Anti-CBL antibody ab59211

2 Images

Overview

Product name	Anti-CBL antibody
Description	Rabbit polyclonal to CBL
Host species	Rabbit
Specificity	ab59211 detects endogenous levels of total CBL protein.
Tested applications	Suitable for: ELISA, IHC-P, WB
Species reactivity	Reacts with: Human Predicted to work with: Mouse, Rat 
Immunogen	Synthetic non-phosphopeptide derived from human CBL around the phosphorylation site of tyrosine 700 (T-E-Y ^P -M-T).
Positive control	IHC-P: Human breast carcinoma tissue. WB: 293 cell extracts.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
Storage buffer	Preservative: 0.02% Sodium Azide Constituents: 50% Glycerol, PBS (without Mg ²⁺ and Ca ²⁺), 150mM Sodium chloride, pH 7.4
Purity	Immunogen affinity purified
Purification notes	ab59211 was affinity purified from rabbit antiserum by affinity chromatography using epitope specific immunogen.
Clonality	Polyclonal
Isotype	IgG


Applications

Our [Abpromise guarantee](#) covers the use of **ab59211** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ELISA		
IHC-P		
WB		
Application notes	<p>ELISA: 1/5000.</p> <p>IHC-Paraffin: 1/50 - 1/100.</p> <p>WB: 1/500 - 1/1000. Detects a band of approximately 110 kDa (predicted molecular weight: 99 kDa).</p> <p>Not yet tested in other applications.</p> <p>Optimal dilutions/concentrations should be determined by the end user.</p>	
Target		
Function	<p>Participates in signal transduction in hematopoietic cells. Adapter protein that functions as a negative regulator of many signaling pathways that start from receptors at the cell surface. Acts as an E3 ubiquitin-protein ligase, which accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and then transfers it to substrates promoting their degradation by the proteasome. Recognizes activated receptor tyrosine kinases, including PDGFA, EGF and CSF1, and terminates signaling.</p>	
Pathway	Protein modification; protein ubiquitination.	
Involvement in disease	<p>Defects in CBL are the cause of Noonan syndrome-like disorder (NSL) [MIM:613563]. NSL is a syndrome characterized by a phenotype reminiscent of Noonan syndrome. Clinical features are highly variable, including facial dysmorphism, short neck, developmental delay, hyperextensible joints and thorax abnormalities with widely spaced nipples. The facial features consist of triangular face with hypertelorism, large low-set ears, ptosis, and flat nasal bridge. Some patients manifest cardiac defects.</p>	
Sequence similarities	<p>Contains 1 Cbl-PTB (Cbl-type phosphotyrosine-binding) domain.</p> <p>Contains 1 RING-type zinc finger.</p> <p>Contains 1 UBA domain.</p>	
Domain	<p>The RING-type zinc finger domain mediates binding to an E2 ubiquitin-conjugating enzyme. The N-terminus is composed of the phosphotyrosine binding (PTB) domain, a short linker region and the RING-type zinc finger. The PTB domain, which is also called TKB (tyrosine kinase binding) domain, is composed of three different subdomains: a four-helix bundle (4H), a calcium-binding EF hand and a divergent SH2 domain.</p>	
Post-translational modifications	<p>Phosphorylated on tyrosine residues by EGFR, SYK, FYN and ZAP70 (By similarity).</p> <p>Phosphorylated on tyrosine residues by INSR.</p>	
Cellular localization	Cytoplasm.	


Images



Peptide - +

Immunohistochemistry (Paraffin-embedded sections)
- CBL antibody (ab59211)

ab59211, at 1/50 - 1/100 dilution, staining CBL in paraffin embedded human breast carcinoma tissue by Immunohistochemistry in the absence or presence of the immunising peptide.



Western blot - CBL antibody (ab59211)

All lanes : Anti-CBL antibody (ab59211) at 1/500 dilution

Lane 1 : 293 cell extract

Lane 2 : 293 cell extract with immunising peptide

Predicted band size: 99 kDa

Observed band size: 110 kDa

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors