

Product datasheet

Anti-JNK1 antibody ab61190

1 Image

Overview

Product name	Anti-JNK1 antibody
Description	Mouse monoclonal to JNK1
Host species	Mouse
Tested applications	Suitable for: WB, ELISA
Species reactivity	Reacts with: Recombinant fragment Predicted to work with: Human
Immunogen	JNK1 tagged recombinant full length protein (Human).
Positive control	Recombinant protein.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
Storage buffer	Preservative: None Constituents: PBS, pH 7.2
Purity	Protein A purified
Clonality	Monoclonal
Isotype	IgG2a

Applications

Our [Abpromise guarantee](#) covers the use of **ab61190** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		
ELISA		

Application notes

ELISA: Use at an assay dependent dilution.

WB: Use at a concentration of 1 - 5 µg/ml. Detects a band of approximately 70 kDa (predicted molecular weight: 48 kDa).

This antibody has only been tested in WB against the recombinant fragment used as immunogen. We have no data on the detection of endogenous protein.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

Target

Function

Responds to activation by environmental stress and pro-inflammatory cytokines by phosphorylating a number of transcription factors, primarily components of AP-1 such as JUN, JDP2 and ATF2 and thus regulates AP-1 transcriptional activity. In T-cells, JNK1 and JNK2 are required for polarized differentiation of T-helper cells into Th1 cells (By similarity).

Phosphorylates heat shock factor protein 4 (HSF4).

JNK1 isoforms display different binding patterns: beta-1 preferentially binds to c-Jun, whereas alpha-1, alpha-2, and beta-2 have a similar low level of binding to both c-Jun or ATF2. However, there is no correlation between binding and phosphorylation, which is achieved at about the same efficiency by all isoforms.

Sequence similarities

Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily.

Contains 1 protein kinase domain.


Domain

The TXY motif contains the threonine and tyrosine residues whose phosphorylation activates the MAP kinases.

Post-translational modifications

Dually phosphorylated on Thr-183 and Tyr-185, which activates the enzyme.

Images



Anti-JNK1 antibody (ab61190) at 1 µg/ml +
recombinant protein at 0.2 µg

Secondary

Goat Anti-Mouse IgG (H&L)-HRP at 1/5000
dilution

Predicted band size: 48 kDa

Observed band size: 70 kDa

Western blot against tagged recombinant
protein immunogen using ab61190 JNK1
antibody. Predicted band size of immunogen
is 48 kDa.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors