

Product datasheet

Anti-Plasminogen antibody ab61387

Overview

Product name	Anti-Plasminogen antibody
Description	Rat monoclonal to Plasminogen
Host species	Rat
Tested applications	Suitable for: WB, ELISA
Species reactivity	Reacts with: Mouse
Immunogen	Full length native purified Mouse Plasminogen

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Preservative: None Constituents: 50% (v/v)Glycerol/water
Purity	Ion Exchange Chromatography
Purification notes	Salt fractionation followed by gel filtration and ion exchange chromatography.
Clonality	Monoclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab61387** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent dilution. Predicted molecular weight: 91 kDa.
ELISA		Use at an assay dependent dilution.

Target

Function	<p>Plasmin dissolves the fibrin of blood clots and acts as a proteolytic factor in a variety of other processes including embryonic development, tissue remodeling, tumor invasion, and inflammation. In ovulation, weakens the walls of the Graafian follicle. It activates the urokinase-type plasminogen activator, collagenases and several complement zymogens, such as C1 and C5. Cleavage of fibronectin and laminin leads to cell detachment and apoptosis. Also cleaves fibrin, thrombospondin and von Willebrand factor. Its role in tissue remodeling and tumor invasion may be modulated by CSPG4. Binds to cells.</p> <p>Angiostatin is an angiogenesis inhibitor that blocks neovascularization and growth of experimental primary and metastatic tumors in vivo.</p>
Tissue specificity	Present in plasma and many other extracellular fluids. It is synthesized in the liver.
Involvement in disease	<p>Defects in PLG are a cause of susceptibility to thrombosis (THR) [MIM:188050]. It is a multifactorial disorder of hemostasis characterized by abnormal platelet aggregation in response to various agents and recurrent thrombi formation.</p> <p>Defects in PLG are the cause of plasminogen deficiency (PLGD) [MIM:217090]. PLGD is characterized by decreased serum plasminogen activity. Two forms of the disorder are distinguished: type 1 deficiency is additionally characterized by decreased plasminogen antigen levels and clinical symptoms, whereas type 2 deficiency, also known as dysplasminogenemia, is characterized by normal, or slightly reduced antigen levels, and absence of clinical manifestations. Plasminogen deficiency type 1 results in markedly impaired extracellular fibrinolysis and chronic mucosal pseudomembranous lesions due to subepithelial fibrin deposition and inflammation. The most common clinical manifestation of type 1 deficiency is ligneous conjunctivitis in which pseudomembranes formation on the palpebral surfaces of the eye progresses to white, yellow-white, or red thick masses with a wood-like consistency that replace the normal mucosa.</p>
Sequence similarities	<p>Belongs to the peptidase S1 family. Plasminogen subfamily.</p> <p>Contains 5 kringle domains.</p> <p>Contains 1 PAN domain.</p> <p>Contains 1 peptidase S1 domain.</p>
Domain	Kringle domains mediate interaction with CSPG4.
Post-translational modifications	<p>N-linked glycan contains N-acetyllactosamine and sialic acid. O-linked glycans consist of Gal-GalNAc disaccharide modified with up to 2 sialic acid residues (microheterogeneity).</p> <p>In the presence of the inhibitor, the activation involves only cleavage after Arg-580, yielding two chains held together by two disulfide bonds. In the absence of the inhibitor, the activation involves additionally the removal of the activation peptide.</p>
Cellular localization	Secreted. Locates to the cell surface where it is proteolytically cleaved to produce the active plasmin. Interaction with HRG tethers it to the cell surface.
Form	Cleaved into the following 5 chains: 1.Plasmin heavy chain A2.Activation peptide3.Angiostatin4.Plasmin heavy chain A, short form5. Plasmin light chain B

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors